

Volume 13, № 47, September 2025

ISSN: 2311 — 1623 (Print)

ISSN: 2311 — 1631 (Online)

<http://www.heart-vdj.com>



International Heart and Vascular Disease Journal

Journal of the Cardioprogress Foundation



Comparative efficacy of pharmacological
and non-pharmacological
management of monomorphic
right ventricular premature
contractions in patients
without structural heart
disease

Features of acute coronary
syndrome associated with
anxiety and depression
disorders

2025 ESC Congress HOT
LINE sessions: key clinical
studies

Editor-in-Chief: **Mekhman Mamedov**

Deputy Editor: **Sergey Kanorsky**

Senior Consulting Editors: **Nathan Wong**

Richard Williams

EDITOR-IN-CHIEF

Mekhman Mamedov, Russia

DEPUTY EDITOR

Sergey Kanorsky, Russia

SENIOR CONSULTING EDITORS

Nathan Wong, USA

Richard Williams, UK

STATISTICAL CONSULTANT

Alexander Deev, Russia

EDITORIAL BOARD

Arabidze G.G., Russia

Vasyuk Yu.A., Russia

Rasulova Z.D., Uzbekistan

Mitchenko E.I., Ukraine

Khirmanov V.N., Russia

Tsinamdzgvrishvili B.V., Georgia

Yakubova L.V., Belarus

Adnan A., Turkey

Dayi H., China

Dusko V., Republika Srpska, Bosnia and Herzegovina

Kazuaki T., Japan

Maciej B., Poland

Poponina T.M., Russia

Pekka P. Finland

Pranas S., Lithuania

Seth J. Baum, USA

Murkarmilov I. L., Kyrgyzstan

EXECUTIVE EDITOR

E.A. Savchuk, Russia

EXECUTIVE TRANSLATOR

G. Ginoyan, Russia

CONTACT DETAILS:

Cardioprogress Foundation and Editorial Office:

Room 213, Building 2, Prospect

Gostinichny 6, Moscow 127106, Russia

Editorial Office tel.: (+7) 965 236 1600

Official website: <http://www.heart-vdj.com>

Editorial correspondence should be sent to:

Mekhman Mamedov, Editor-in-Chief,

editor.ihvdj@gmail.com

Articles for publication should be sent to:

Anna Artyeva, Associate Editor,

submissions.ihvdj@gmail.com

© International Heart and Vascular Disease Journal is an official publication of the Cardioprogress Foundation

Printed in Russia

The Journal is in the List of the leading scientific journals and publications of the Supreme Examination Board (VAK)

Complete versions of all issues are published:

www.elibrary.ru, www.cyberleninka.ru

International Heart and Vascular Disease Journal

Journal of the «Cardioprogress» Foundation

Volume 13, № 47, September 2025

Contents

Editor's welcome	2
Review of international medical news	3

LEADING ARTICLE

Olesin A. I., Konstantinova I. V., Tyuteleva N. N., Zueva I.

Comparative efficacy of pharmacological and non-pharmacological management of monomorphic right ventricular premature contractions in patients without structural heart disease	4
--	---

ORIGINAL ARTICLES

Obedkova N.Yu., Mal G.S., Tsukanov A.V.

An innovative approach to assessing residual risk using advanced glycation end product autofluorescence in patients with coronary heart disease based on various lipid-lowering therapy regimens	11
---	----

Kovalenko E. V., Belaya O. L., Markova L. I., Evdokimov V. V.

Prognostic value of selected biomarkers in comorbid patients with chronic heart failure after COVID-19	17
---	----

Yupatova M. I., Nikitina N. M., Gafurova L. R.

Clinical significance of 24-hour blood pressure monitoring in patients with ankylosing spondylitis	26
---	----

REVIEW ARTICLE

Poponina T. M., Repin A. N., Poponina Y. S., Gunderina K. I.

Features of acute coronary syndrome associated with anxiety and depression disorders	31
---	----

Panakhova D. Z., Mirzabekova K. N.

Modern trends in hypolipidemic therapy	37
---	----

REPORT

2025 ESC Congress HOT LINE sessions: key clinical studies	48
--	----

Author guidelines	52
--------------------------------	----



Editor's Welcome

Dear colleagues!

We present to your attention the forty-seventh issue of the International Heart and Vascular Disease Journal, which features leading, original, and review articles, as well as a report on key clinical studies presented during the HOT LINE sessions of the 2025 Congress of the European Society of Cardiology.

In the "Leading Article" section, an original study is presented that evaluates the effectiveness of pharmacological therapy and radiofrequency catheter ablation for the treatment of monomorphic right ventricular extrasystole, depending on the presence of predictors of arrhythmogenic cardiomyopathy in patients without structural heart abnormalities. The study included 452 patients, with a follow-up period of up to 5 years. According to the authors, among patients without structural heart abnormalities who had early monotypic right ventricular extrasystole without predictors of arrhythmogenic cardiomyopathy, Class I antiarrhythmic drugs were found to be the most effective. Meanwhile, in patients with monomorphic right ventricular extrasystole and identified predictors of arrhythmogenic cardiomyopathy, radiofrequency catheter ablation proved to be the most effective treatment approach.

The "Original Articles" section features three studies. The first article compares the levels of asymmetric dimethylarginine in patients with coronary heart disease (CHD) with and without metabolically associated fatty liver disease. The study included 50 patients diagnosed with CHD, divided into two equal groups: 25 with metabolically associated fatty liver disease and 25 without it. The results indicate the need for a personalized approach to treating patients with CHD and metabolically associated fatty liver disease, focusing on correcting metabolic disorders to reduce asymmetric dimethylarginine levels. The second article demonstrates that the use of N-terminal pro-B-type natriuretic peptide, cystatin C, and galectin-3 as part of a multifactorial model allows for more effective prediction of heart failure decompensation and cardiovascular mortality in post-COVID-19 patients with chronic heart failure with preserved ejection fraction, type 2 diabetes mellitus, and chronic kidney disease. The third study shows that arterial hypertension (AH) in patients with spondyloarthritis may present subclinically. About half of the patients who denied having AH were found to have masked hypertension, and among them, half had an abnormal circadian blood pressure pattern. The authors conclude that 24-hour blood pressure monitoring is necessary for the early detection of hypertension in patients with ankylosing spondylitis.

The "Review Articles" section includes two papers. The first article states that psychological and pharmacological interventions can have beneficial effects and should be considered for patients with acute coronary syndrome who experience depression, anxiety, and stress. Before discharge, all patients are recommended to be screened for mental disorders using validated questionnaires and referred for psychiatric consultation if necessary. The second article presents data on new lipid-lowering drugs. Bempedoic acid, an ATP-citrate lyase inhibitor, is emerging as a promising treatment option for patients with a history of statin intolerance. Icosapent ethyl, a high-dose ethyl ester of eicosapentaenoic acid, has demonstrated favorable effects on cardiovascular health.

As is traditional, the issue concludes with a brief report summarizing key clinical trials presented at the HOT LINE sessions of the 2025 European Society of Cardiology Congress.

We invite everybody to collaborate with the journal. Our team is waiting for your original papers, review articles, discussions, and opinions about problems, treatment and prophylaxis recommendations.

Mekhman N. Mamedov

Editor-in-Chief

President of the "Cardioprogress" Foundation

Review of international medical news

Scientists from Scotland conducted a study showing that in patients who had suffered a myocardial infarction (MI), a three-month course of dual antiplatelet therapy produced comparable outcomes to traditional 12-month treatment. At the same time, it was associated with reduced mortality and a lower risk of bleeding.

The analysis revealed that all-cause mortality was 2.7% in the three-month treatment group and 3.4% in the 12-month group. There were no significant differences in cardiovascular mortality or non-fatal MI rates.

The authors concluded that three-month dual antiplatelet therapy after MI may be a safe alternative to the standard 12-month regimen. This approach reduces bleeding risk without compromising efficacy in preventing recurrences, which may be taken into account in future clinical guidelines.

According to ESC data

Experts have issued a scientific statement on the impact of psychological distress on outcomes in patients after myocardial infarction. The document reviews the mechanisms of these disorders, their connection with cardiovascular risk, and provides recommendations for diagnosis and management.

The analysis showed that depression, anxiety, post-traumatic stress disorder, and chronic stress after myocardial infarction significantly increase the risk of recurrent cardiovascular events and mortality. According to the SWEDEHEART registry, 21% of patients experienced persistent emotional distress, which increased the risk of cardiovascular death by 46%.

Numerous studies indicate that post-infarction depression doubles the likelihood of recurrent cardiovascular events.

According to the Circulation journal

Specialists from an international research group conducted a systematic meta-analysis of tumor necrosis factor- α (TNF- α) inhibitor therapy in patients with rheumatoid arthritis.

The analysis showed that TNF- α inhibitor use was associated with a significant increase in the risk of serious infections (odds ratio = 1.63). However, this therapy reduced the risk of cardiovascular events by 40% and myocardial infarction by 25%.

The authors concluded that while TNF- α inhibitor therapy in rheumatoid arthritis patients carries a high risk of infections, but may reduce the likelihood of cardiovascular events and improve metabolic syndrome parameters.

According to The Lancet journal

Researchers from France found that prescribing beta-blockers after myocardial infarction does not affect mortality, recurrence, or hospitalization rates in patients with heart failure and preserved ejection fraction.

The study included 8,500 patients with heart failure and a left ventricular ejection fraction above 40% who had experienced myocardial infarction. During 3.7 years of follow-up, adverse outcomes were observed in 316 participants in the beta-blocker group (22.5 events per 1,000 patient-years) and in 307 participants not receiving beta-blockers (21.7 events per 1,000 patient-years).

The researchers noted that when prescribing these medications to such patients, the obtained results should be taken into consideration.

According to the NEJM

Researchers from the United States analyzed the role of lipoprotein(a) as a biomarker for extracoronary atherosclerotic complications.

The analysis showed that an increase in lipoprotein(a) levels by 75 nmol/L was associated with an 18% higher risk of peripheral artery disease and a 17% higher risk of carotid artery stenosis. In patients with pre-existing peripheral artery disease, elevated lipoprotein(a) levels increased the risk of serious limb-related adverse events by 57%.

Thus, this study demonstrates that lipoprotein(a) is an important independent marker for the development and progression of extracoronary atherosclerotic disease

According to the Circulation journal

Scientists from Korea conducted a study involving 241,924 adults, which found that increasing the cumulative CVH score (Life's Essential 8) by 100 points per year between ages 30 and 40 was associated with a 34% lower risk of cardiovascular events and a 35% lower risk of kidney events in middle age.

The results suggest that maintaining a high cumulative CVH score (points $\times$ years) may serve as a valuable target for primary prevention, as achieving and maintaining good cardiovascular health earlier in life is associated with a reduced risk of premature cardiovascular and renal events in the 30–40 year age range.

According to the JAMA Cardiology

Comparative efficacy of pharmacological and non-pharmacological management of monomorphic right ventricular premature contractions in patients without structural heart disease

Olesin A.I., Konstantinova I.V., Tyuteleva N.N., Zueva I.S.

Federal State Budget Institution of Higher Education "North-Western State Medical University Named After I.I Mechnikov" Under the Ministry of Public Health of the Russian Federation, Saint-Petersburg, Russia.

AUTHORS

Alexander I. Olesin, MD, PhD, Professor of the Department of Hospital Therapy and Cardiology named after M. S. Kushakovskiy, Federal State Budget Institution of Higher Education "North-Western State Medical University Named After I.I Mechnikov" Under the Ministry of Public Health of the Russian Federation, Saint-Petersburg, Russia. ORCID: 0000-0001-7827-1052

Irina V. Konstantinova, MD, PhD, Federal State Budget Institution of Higher Education "North-Western State Medical University Named After I.I Mechnikov" Under the Ministry of Public Health of the Russian Federation, Saint-Petersburg, Russia. ORCID: 0000-0003-3350-3088

Natalya N. Tyuteleva, MD, PhD, Assistant Lecturer of the Department of Hospital Therapy and Cardiology named after M. S. Kushakovskiy, Federal State Budget Institution of Higher Education "North-Western State Medical University Named After I.I Mechnikov" Under the Ministry of Public Health of the Russian Federation, Saint-Petersburg, Russia. ORCID: 0000-0002-9083-8944

Iuliia S. Zueva, MD, PhD, Assistant Lecturer of the Department of Hospital Therapy and Cardiology named after M. S. Kushakovskiy, Federal State Budget Institution of Higher Education "North-Western State Medical University Named After I.I Mechnikov" Under the Ministry of Public Health of the Russian Federation, Saint-Petersburg, Russia. ORCID: 0000-0002-1685-6300

To date, antiarrhythmic medication and radiofrequency ablation (RFA) are the primary methods of right ventricular premature contractions (PVCs) management in patients without structural heart disease (SHD).

Aim of the study. The aim of the study was to evaluate the efficacy of pharmacological and non-pharmacological (RFA) strategies of monomorphic right ventricular PVCs management, depending on the identification of

arrhythmogenic cardiomyopathy (ACM) predictors in patients without SHD.

Methods. In this study, 452 patients with high frequency of PVCs (mean 23529 ± 561 PVCs per day) and without SHD aged 16-39 were observed. In addition to the basic parameters, such ACM predictors as the QRS complex maximum intrinsicoid deflection index, QRS complex duration, etc. were electrocardiographically assessed. The observation period was 5 years.

Results. In 95.13% of patients the early monotypic right ventricular PVCs without ACM predictors were diagnosed. No cardiovascular diseases developed in these patients, and Class I antiarrhythmic drugs were the most effective at suppressing ventricular ectopy. In this group, 15.71% of the patients declined the long-term antiarrhythmic therapy or RFA. The rest of the group continued pharmacologic therapy. In 95.28% of patients with the early monotypic right ventricular PVC without registered ACM predictors, the duration of antiarrhythmic therapy, according to the "therapeutic sufficiency" principle, was 11.1 ± 0.3 months from the beginning of the period of observation. This duration highly correlated ($r = 0.89$) with the observation period of patients receiving cardioprotective therapy. In 4.87% of patients with registered right ventricular PVC with ACM predictors, no effect of antiarrhythmic therapy was observed and/or clinically significant heart failure

developed. These patients underwent RFA for ventricular ectopy management. In all patients of this subgroup RFA induced remission and in 85.71% of them remained free of PVC recurrence throughout the entire follow-up period.

Conclusion. Patients with the early monotypic right ventricular PVCs, without SHD and any ACM predictors treatment with Class I antiarrhythmics was the most effective, while in patients with monomorphic right ventricular PVCs and ACM predictors, RFA was the effective therapeutic option.

Keywords: Premature ventricular contractions, antiarrhythmic therapy personalization, pharmacologic treatment, radiofrequency ablation.

Conflict of interests: none declared.

Accepted: 19.06.2025

Received: 01.08.2025



For citation: Olesin A.I., Konstantinova I.V., Tyuteleva N.N., Zueva I.S. Comparative efficacy of pharmacological and non-pharmacological management of monomorphic right ventricular premature contractions in patients without structural heart disease. 2025. 13(47):4-10. DOI 10.24412/2311-1623-2025-47-5-12

Introduction

In patients without structural heart disease (SHD), in most cases, good prognosis of monomorphic right ventricular premature contractions (PVCs) is observed, except for the development of myocardial contractility depression in high frequency ventricular ectopy [1, 2]. Despite the use of various cardiac mapping methods, which assist with the definition of the source of premature ventricular complexes, they might be ineffective for the verification of PVCs mechanism in patients without SHD. In particular, these methods can not help to differentiate between trigger activity and re-entry mechanism [3, 4].

Prior clinical and experimental studies have shown that in most cases, early monomorphic right ventricular PVCs in patients without SHD and in the absence of electrocardiographic (ECG) predictors of arrhythmogenic cardiomyopathy, such as the duration of

the ectopic QRS complex and the sinus rhythm QRS complex, their ratio, the measurement of the maximum intrinsicoid deflection index of the ectopic complex, and the variability of the corrected pre-ectopic interval assessed by its linear deviation (LD), etc., is caused by a mechanism of early afterdepolarization (EAD) and follows a predominantly benign course. Furthermore, prospective follow-up of these patients revealed no subsequent cardiovascular pathology [5].

The available literature lacks data on the comparative efficacy of pharmacological therapy versus non-pharmacological management, specifically radiofrequency catheter ablation (RFA), as well as on the natural history of early monomorphic right ventricular PVCs in patients without SHD. This is specifically for cases predominantly caused by an EAD mechanism, for instance, when antiarrhythmic drug therapy is declined to eliminate the ventricular ectopy.

Aim of the study

The aim of the study was to evaluate the efficacy of pharmacological and non-pharmacological (RFA) strategies of monomorphic right ventricular PVCs management, depending on the identification of arrhythmogenic cardiomyopathy (ACM) predictors in patients without SHD.

Methods

A total of 452 patients without SHD, aged 16 to 39 years (mean 22.1 ± 0.1 years), with monomorphic right ventricular PVCs of IV-V gradation according to the M. Ryan classification [6], were observed.

The inclusion criteria were: documented sinus rhythm; absence of SHD, confirmed by ruling out cardiac and extracardiac diseases, electrolyte imbalances, use of medications and/or toxic products (primarily diuretics, oral contraceptives, alcohol abuse, etc.) that directly or indirectly lead to the development of ventricular arrhythmias; confirmation of the absence of SHD using various stress tests, invasive and non-invasive coronary angiography, echocardiography, and contrast-enhanced cardiac magnetic resonance imaging; presence of subjective palpitations; left ventricular ejection fraction $\geq 54\%$; and provision of informed consent for participation in the study and treatment (as previously described) [5, 7].

All patients underwent general clinical examination, 24-72-hour Holter ECG monitoring, and echocardiography using a Hitachi EUB-5500 device according to standard protocols. The calculation of hemodynamic parameters, determination of the topographic localization of the premature ventricular complexes, comparison of the pre-ectopic interval (PEI) of the PVC to the QT interval of the sinus rhythm (QTsr), determination of the linear deviation (LD) of the corrected PEI (PEIcor), measurement of the QRS duration of the PVC (QRSpvc) and sinus rhythm (QRSsr), their ratio (QRSpvc/QRSsr), and the maximum intrinsicoid deflection time (MIDpvc) of the right ventricular extrasystole were performed as previously described [5, 9].

Pathological values for MIDpvc, QRSpvc duration, and the QRSpvc/QRSsr ratio were defined as ≥ 0.5 units, ≥ 150 ms, and ≥ 1.5 units, respectively [2, 5]. Monotopic premature ventricular complexes were defined by an LD of the corrected PEI (PEIcor LD) of < 10 ms [3].

The selection of antiarrhythmic therapy for ventricular ectopy was based on 24-hour Holter ECG

monitoring data. This included the identification of potentially effective Class I-III antiarrhythmic drugs, which were used in medium therapeutic doses [3].

The treatment protocol was initiated with Class II agents. If these were ineffective, Class I or III drugs were used. Amiodarone was employed as a last resort, if necessary. The scope and nature of antiarrhythmic drug therapy, as well as the criteria for treatment efficacy for ventricular ectopy, have been described previously [5, 6].

Pharmacological therapy followed the principle of «therapeutic sufficiency» This meant that antiarrhythmic drugs were withdrawn every 3-6 months. If PVCs recurred, the previous antiarrhythmic pharmacotherapy was reinstated. If no premature ventricular complexes were recorded, the treatment was discontinued [5, 8].

In cases where mono- or combined antiarrhythmic therapy was ineffective, or if depression of myocardial contractility developed leading to a refusal of long-term antiarrhythmic drug use, RFA became the treatment of choice for PVCs in patients without SHD [2, 3, 5].

The duration of patient follow-up after inclusion in the study was 5 years. All assessments, including patient status evaluation and ECG recording, were performed once a month. 24-hour Holter ECG monitoring was conducted no less than once every 3-4 months.

Statistical analysis

For the statistical analysis of the obtained data, the following methods were used: means and standard errors of the mean ($M \pm m$), the 95% confidence interval for the means, Student's t-test, and the χ^2 test. A p-value of less than 0.05 ($p < 0.05$) was considered statistically significant. The normality of distribution for quantitative variables was assessed using the Kolmogorov-Smirnov and Gaussian tests. Linear pairwise correlation (Pearson's r) and rank correlation (Spearman's r) were employed. The association between quantitative and qualitative binary variables was evaluated using logistic regression analysis with the calculation of odds ratios (OR).

Results

Among the patients without SHD included in the study, the number of PVCs over 24 hours ranged from 11,256 to 39,798 (mean $23,529 \pm 561$ PVCs per day). Furthermore, 384 patients (84.96%) had ventricular ectopy accounting for 15% or more of all ventricular

complexes, and 98 patients (21.68%) had non-sustained ventricular tachycardia.

The most effective drug for suppressing PVCs was etacizin in 356 patients (78.76%), allapinin in 58 (12.83%), propafenone in 23 (5.09%), Class II agents in 9 (2.00%), and sotalol in the remaining 6 patients (1.32%).

Seventy-one patients (15.71%) declined long-term antiarrhythmic drug therapy for various reasons, including the development of headaches, dizziness, diplopia, nausea, constipation or diarrhea, allergic reactions, financial constraints, etc. These patients received cardioprotective therapy, which included cardiometabolic agents such as potassium supplements and polyunsaturated fatty acids. Among them, 4 patients (5.63%) underwent RFA 6-8 months after the start of observation to eliminate ventricular ectopy due to the development of impaired pump function, which manifested as clinically overt heart failure. The remaining 381 patients (84.29%) continued antiarrhythmic medication. Of these, 18 patients (4.72%) underwent RFA 7-14 months (mean 10.8 ± 0.9 months) after initiating therapy due to the ineffectiveness of pharmacological antiarrhythmic treatment, which included the use of Class I-III drugs and their combinations.

All patients were divided into two groups. Group I consisted of 430 patients (95.13%) who used drug therapy to manage PVCs. The remaining patients were included in Group II and underwent RFA to eliminate ventricular ectopy. Group I was further divided into two subgroups. Subgroup Ia included 67 patients (15.58%) who were treated for PVCs with cardiomet-

abolic agents only; this subgroup was used to study the «natural history» of right ventricular ectopy in patients without SHD. Subgroup Ib included 363 patients (84.42%) who received pharmacological antiarrhythmic therapy to eliminate PVCs.

All patients in subgroups Ia and Ib exhibited early monomorphic premature ventricular contractions (PVCs) without pathological values of ACM predictors. No statistically significant difference in the studied parameters was found between these subgroups (Table 1).

In Group II, all parameters reflecting ACM predictors for PVCs were significantly higher compared to subgroups Ia and Ib. No significant changes in the other studied parameters were observed between subgroups Ia and Ib when compared to Group II.

A positive clinical effect was observed in 32 patients (47.76%) of subgroup Ia within 3-9 months (mean 6.2 ± 0.5 months), and in the remaining patients within 10-23 months (mean 16.3 ± 0.8 months). In all patients of this subgroup, the PVC burden decreased by an average of 92% compared to baseline.

In 178 patients (49.04%) of subgroup Ib, the duration of antiarrhythmic therapy, guided by the principle of «therapeutic sufficiency», ranged from 4 to 10 months (mean 6.7 ± 0.2 months) from the start of observation. For the remaining patients, it ranged from 11 to 21 months (mean 16.7 ± 0.5 months). This duration strongly correlated with the observation period of patients in subgroup Ia ($r = 0.89$).

In all patients of Group II, the PVCs were eliminated immediately after RFA. A positive effect from RFA was maintained throughout the entire follow-up

Table 1. Electrocardiographic parameters of PVCs in subgroups Ia, Ib, and group II patients

Patient groups Parameters	Subgroup Ia n = 67	Subgroup Ib n = 363	Group II n = 22
PEI _{pvc} /QTSR, ratio	1.26 ± 0.01 (1.18-1.36)	1.25 ± 0.01 (1.14-1.36)	1.75 ± 0.12 (1.22-2.32)*
LD of cor. PEI pvc, ms	8.62 ± 0.08 (7.97-9.26)	8.61 ± 0.01 (8.12-9.14)	82.35 ± 10.51 (39-126)*
MID _{pvc} , ratio (PV ≥ 0.5)	0.41 ± 0.01 (0.31-0.52)	0.42 ± 0.01 (0.32-0.53)	0.58 ± 0.01 (0.44-0.71)* +
QRS duration of PVC, ms (PV ≥ 150 ms)	137 ± 1 (125-152)	138 ± 1 (122-154)	179 ± 5 (145-212)* +
QRS _{pvc} /QRS _{sr} , ratio (PV ≥ 1.5)	1.38 ± 0.01 (1.24-1.53)	1.39 ± 0.01 (1.22-1.56)	1.88 ± 0.07 (1.46-2.23)* +
Number of PVCs per 24 hours	23787 ± 1235 (11657-35875)	23423 ± 531 (14856-31987)	24787 ± 2456 (14678-36784)

Note: * — significant difference compared to subgroup Ia, ♦ — significant difference compared to subgroup Ib, + — significant difference compared to pathological values ($p < 0.05$).

period in 16 patients (72.73%) of this group. In 6 patients (27.27%), PVCs recurred 2 years after the ablation and were successfully eliminated with a repeat RFA procedure; the morphology of the recurrent premature ventricular complexes was similar to the original. Endomyocardial biopsy was performed in 8 patients (36.36%) of this group, revealing myocardial inflammatory fibrosis (myocarditic cardiosclerosis).

The positive effect of RFA highly correlated with the following ACM predictors: MID of the PVC ≥ 0.5 units (OR = 15.4), a PVC QRS complex duration ≥ 150 ms (OR = 12.9), and a ratio of PVC QRS to sinus rhythm QRS duration ≥ 1.5 units (OR = 9.6).

No cardiovascular diseases were detected in any patient of Group I throughout the entire observation period.

Discussion

It is currently known that frequent monomorphic right ventricular PVCs are detected in 3% to 5% of young patients (typically under 40 years of age) without SHD, and in most cases, are typically benign [2, 3]. In patients without SHD, the indications for antiarrhythmic therapy for monomorphic PVCs are: the presence of subjective symptoms, or the detection of ventricular ectopy accounting for 15% or more of total heartbeats. If patients decline pharmacologic antiarrhythmic therapy or if it proves ineffective, RFA becomes the treatment of choice for eliminating the premature ventricular complexes [1-3].

A total of 452 patients without SHD, aged 16 to 39 years, with frequent monomorphic PVCs (mean $23,529 \pm 561$ PVCs per 24 hours) were observed. Early monomorphic right ventricular PVCs without predictors of ACM were recorded in 95.13% of them.

Clinical and experimental studies have shown that early monomorphic right ventricular PVCs in patients without SHD and without ACM predictors is most often caused by a mechanism of early afterdepolarization (EAD) [5].

EADs are caused by an increase in the membrane potential of cardiomyocytes to a level between -40 mV and -10 mV. This can result from, for example, inhibition or delayed inactivation of the sodium channel system, leading to the emergence of spontaneous action potentials [10, 11]. This phenomenon can be induced experimentally by various toxic agents, such as aconitine, and in humans, it may result from «oxidative stress» or other local metabolic disturbances affecting

cardiomyocyte function [10, 11]. For suppressing ventricular ectopy caused by the EAD mechanism, Class I antiarrhythmic drugs are the most effective [12].

Similar findings were obtained in the present study: etacizin was the most effective drug in 78.76% of patients, allapinin in 12.83%, and propafenone in 5.09%.

For various reasons, 15.71% of patients declined long-term antiarrhythmic pharmacotherapy, while the remaining patients continued treatment with antiarrhythmic drugs.

In patients without SHD presenting with early monomorphic right ventricular PVCs, the rationale for antiarrhythmic therapy extends beyond preventing «arrhythmia-induced cardiomyopathy.» The fact that frequent ventricular ectopy itself may induce disturbances in myocardial metabolism—potentially through increased sympathetic activity, ventricular dyssynchrony, and other factors leading to stabilization of the ectopic focus, also serves as a basis for treatment [13-15].

In patients without SHD receiving cardioprotective therapy, a positive clinical effect was observed after an average of 11.8 ± 0.7 months: the PVC burden decreased by an average of 92% compared to baseline. The duration of pharmacological antiarrhythmic therapy in patients without SHD, guided by the principle of «therapeutic sufficiency,» averaged 11.1 ± 0.3 months from the start of observation. This duration strongly correlated with the observation period of patients receiving cardioprotective therapy ($r = 0.89$).

In patients without SHD, the occurrence of early monomorphic right ventricular PVCs without ACM predictors, typically caused by trigger mechanisms such as EAD, reflects a reversible dysfunction of cardiomyocytes [1, 5, 6]. This is supported by the results of the present study: no cardiovascular pathology was detected in these patients throughout the entire follow-up period.

In 4.87% of patients without SHD who had monomorphic PVCs with ACM predictors and who declined antiarrhythmic drug therapy, RFA was performed 6-14 months after the start of observation to eliminate ventricular ectopy due to the development of clinical heart failure and/or inefficacy of pharmacological antiarrhythmic therapy.

The immediate success rate of RFA, used, for example, when long-term antiarrhythmic drug therapy is declined or ineffective—is 80-90% [1-3], and according to some data, reaches nearly 100% [14,

15]. Similar results were obtained in the present study.

The development of monomorphic PVCs with ACM predictors in patients without SHD is likely caused by the formation of small focal areas of fibrosis or cardiosclerosis, which are usually not detected by conventional diagnostic methods. These areas may facilitate the generation of premature ventricular complexes, possibly through the induction of an ectopic focus or a re-entry mechanism [10–12]. Furthermore, previous clinical and experimental studies have shown that the identification of ACM predictors in patients without SHD with PVCs indicates an organic cause of the arrhythmia and is a risk factor for the development of cardiovascular pathology [5].

This notion is indirectly confirmed by the results of the present study. A positive effect from RFA was maintained throughout the entire follow-up period in 72.73% of patients; this outcome highly correlated with the presence of ACM predictors. An endomyocardial biopsy was performed in 36.36% of these patients, revealing myocardial fibrosis consistent with myocarditic cardiosclerosis. In 27.27% of patients, PVCs recurred 2 years after the ablation and were successfully eliminated with a repeat RFA procedure; the morphology of the recurrent premature ventricular complexes was similar to the original.

References

- Scorza R, Jonsson M, Corander JM et al. Prognostic impact of morphology and duration of premature ventricular contractions in a population without structural heart disease. *Ann Noninvasive Electrocardiol.* 2023;28(4):e13067. DOI: 10.1111/anec.13067
- Lavallaz JF, Mézier J, Mertz L et al. Risk factors for the development of premature ventricular complex-induced cardiomyopathy: a systematic review and meta-analysis. *J. Interv. Card. Electrophysiol.* 2023;66(5):1145-1163. DOI: 10.1007/s10840-022-01421
- Zeppenfeld K., Tfelt-Hansen J., De Riva M. et al. 2022 ESC Guidelines for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death Developed by the task force for the management of patients with ventricular arrhythmias and the prevention of sudden cardiac death of the European Society of Cardiology (ESC) Endorsed by the Association for European Paediatric and Congenital Cardiology (AEPC). *Europ. Heart J.* 2022;43(40):3997-4126. DOI: 10.1093/eurheartj/ehac262
- Muser D, Tritto M, Mariani MV et al. Diagnosis and Treatment of Idiopathic Premature Ventricular Contractions: A Stepwise Approach Based on the Site of Origin. *Diagnostics*, 2021; 11(10):1840. DOI: 10.3390/diagnostics11101840
- Olesin AI, Konstantinova IV, Tyuteleva NN et al. Clinical evaluation to identify the predictors of arrhythmogenic cardiomyopathy in patients with ventricular extrasystoles without structural heart changes. Clinical and experimental study. *International Heart and Vascular Disease Journal.* 2023. 11(40):28-38. Russian. DOI:10.24412/2311-1623-2023- 40-28-38
- Libby P, Bonow RO, Mann DL et al. *Braunwald's Heart Disease. A textbook of cardiovascular medicine.* 12th ed., Elsevier Science, 2021. 2032 p.
- Gorenk B, Fisher JD, Kudaiberdieva G et al. Premature ventricular complexes: diagnostic and therapeutic considerations in clinical practice: A state-of-the-art review by the American College of Cardiology Electrophysiology Council. *Journal of Interventional Cardiac Electrophysiology*, 2020;57(1):5–26. DOI: 10.1007/s10840-019-00655-3

Conclusion

In 94.37% of patients without SHD who declined antiarrhythmic drug therapy for early monomorphic right ventricular PVCs, a positive clinical effect was observed after an average of 11.8 ± 0.7 months of cardioprotective therapy. In these patients, the burden of premature ventricular complexes decreased by an average of 92% compared to baseline.

In 95.28% of patients without SHD, the duration of pharmacological antiarrhythmic therapy for early monomorphic right ventricular PVCs, guided by the principle of «therapeutic sufficiency,» highly correlated with the observation period of patients receiving cardioprotective therapy ($r = 0.89$).

The presence of monomorphic PVCs with predictors of ACM in patients without SHD is a predictor of ineffective pharmacological antiarrhythmic therapy and/or the development of clinically overt heart failure. For these patients, RFA is the treatment of choice for eliminating ventricular ectopy. Following this procedure, premature ventricular complexes were eliminated and did not recur throughout the entire follow-up period in 72.73% of patients.

Conflict of interests: none declared.

- 10 Olesin A.I., Konstantinova I.V., Tyuteleva N.N., Zueva I.S.
Comparative efficacy of pharmacological and non-pharmacological management...
DOI: 10.24412/2311-1623-2025-47-5-12
-
8. Tsaregorodtsev DA, Sokolov AV, Vasyukov SS et al. Treatment for ventricular arrhythmias in the absence of structural heart disease: from guidelines to clinical practice. *Terapevticheskii Arkhiv*. 2017;89(12):157-164. Russian. DOI: 10.17116/ter-arkh20178912157-164
 9. Yamada T Twelve-lead electrocardiographic localization of idiopathic premature ventricular contraction origins. *J. Cardiovasc. Electrophysiol*. 2019;30(11):2603-2617. DOI: 10.1111/jce.14152
 10. Callans DJ Josephson`s Clinical Cardiac Electrophysiology: Techniques and Interpretations. 6 ed. Lippincott Williams Wilkins, 2020. 850 p.
 11. Welcome MOW, Dogo D, Mastorakis NE Cellular mechanisms and molecular pathways linking bitter taste receptor signalling to cardiac inflammation, oxidative stress, arrhythmia and contractile dysfunction in heart diseases. *Review Inflammopharmacology*. 2023;31(1):89-117. DOI: 10.1007/s10787-022-01086-9
 12. Joukar S A comparative review on heart ion channels, action potentials and electrocardiogram in rodents and human: extrapolation of experimental insights to clinic. *Lab. Anim. Res*. 2021;37(1):25-33. DOI: 10.1186/s42826-021-00102-3
 13. Kantharia BK, Shah AN Are antiarrhythmic agents indicated in premature ventricular complex-induced cardiomyopathy and when? *J Cardiovasc Electrophysiol*. 2024;35(3):574-582. DOI: 10.1111/jce.16055.
 14. Shen X, Zhu X, Zuo L, Liu X, Qin M Mechanisms and Risk Factors for Premature Ventricular Contraction Induced Cardiomyopathy. *Rev. Cardiovasc. Med*. 2023;24(12):353-365. DOI: 10.31083/j.rcm2412353
 15. Pundi K, Marcus MG Predictors and possible mechanisms of premature ventricular contraction induced cardiomyopathy. *J Cardiovasc Electrophysiol*. 2024;35(3):569-573. DOI: 10.1111/jce.1607

The role of metabolic disorders in the formation of asymmetric dimethylarginine levels in patients with coronary heart disease

Nurillaeva N.M.¹, Makhkamova M.M.², Srojedinova N.Z.²

¹ Tashkent State Medical University, Tashkent, Uzbekistan.

² Republican Specialized Scientific and Practical Medical Center of Cardiology, Tashkent, Uzbekistan.

AUTHORS

Nargiza M. Nurillaeva, PhD, MD, Professor, Head of the Department of Internal Diseases in Family Medicine No. 1 with the basics of preventive medicine, Tashkent State Medical University, Tashkent, Uzbekistan. ORCID: 0000-0001-5520-675X

Munisakhon M. Makhkamova, Doctoral student of the Department of Internal Diseases in Family Medicine No. 1 with the basics of preventive medicine, Tashkent State Medical University, Tashkent, Uzbekistan. ORCID: 0000-0002-2658-6148

Nigora Z. Srojedinova, PhD, MD, Head of the Laboratory of Cardiometabolic and Metabolic Disorders, Republican Specialized Scientific and Practical Medical Center of Cardiology, Tashkent, Uzbekistan. ORCID: 0000-0002-4740-2790

Aim of the study. The aim of the study was to compare levels of asymmetric dimethylarginine (ADMA) in patients with coronary heart disease (CHD) in the presence or absence of metabolic dysfunction-associated steatotic liver disease (MASLD) and assess the possible association between ADMA and metabolic disorders.

Methods. The study included 50 patients with CHD, divided into two equal groups: 25 patients with MASLD and 25 patients without MASLD. The CHD was diagnosed based on clinical, electrocardiographic, and coronarographic data. MASLD was diagnosed using liver ultrasound. ADMA levels were assessed using enzyme-linked immunosorbent assay (ELISA) of the blood serum sample.

Results. The mean age in the MASLD group was 54.2 years, and in the group without MASLD, it was 53.8 years. The mean body mass index (BMI) of patients with MASLD was significantly higher (32.8 kg/m²) than that of patients without MASLD (27.1 kg/m², $p < 0.01$). Arterial hypertension was present in 80 % of MASLD patients and 48 % of patients without MASLD ($p < 0.05$). Type 2 diabetes mellitus (T2DM) was diagnosed in 52 % of MASLD patients and 20 % of patients without MASLD ($p < 0.05$). ADMA levels in patients with MASLD were significantly higher (165.85 ± 24.27 ng/mL) than in patients without MASLD (129.44 ± 20.22 ng/mL, $p < 0.001$). Moreover, the presence of three or more metabolic criteria (e.g., obesity, arterial hyper-

tension, and diabetes mellitus) was associated with even higher ADMA levels (190 ± 20 ng/mL).

Conclusion. The obtained data confirm the necessity of a personalized approach to treating patients with CHD and MASLD, focusing on the correction of metabolic disorders to reduce ADMA levels.

Keywords: coronary heart disease, metabolic dysfunction-associated steatotic liver disease, asymmetric dimethylarginine, endothelial dysfunction, insulin resistance

Conflict of interest: none declared.

Accepted: 29.06.2025

Received: 06.08.2025



For citation: Nurillaeva N.M., Makhkamova M.M., Srojidinova N.Z. The role of metabolic disorders in the formation of asymmetric dimethylarginine levels in patients with coronary heart disease. *International Journal of Heart and Vascular Diseases*. 2025. 13(47):11-16. DOI: 10.24412/2311-1623-2025-47-13-20

Introduction

Coronary heart disease (CHD) remains one of the leading causes of death and disability worldwide [1]. According to the World Health Organization (WHO), cardiovascular diseases (CVDs), including CHD, account for approximately 17.9 million people yearly, making up to 32% of all deaths. CHD alone is responsible for more than 7 million deaths each year¹. The European Society of Cardiology reports that CHD is the primary cause of death in Europe [2], accounting for 38% of all CVD-related deaths.

CVDs continue to be a leading cause of mortality and disability in Uzbekistan. According to the Ministry of Health of the Republic of Uzbekistan, circulatory system diseases account for more than 50% of total mortality². Recent WHO data (Global Health Estimates, 2024 revision) indicate that the age-standardized mortality rate from all CVDs in Uzbekistan in 2021 was ≈ 780 – 800 per 100,000 population (both sexes). This is approximately 3.5 times higher than the average rate for the WHO European Region (≈ 220 per 100,000) and nearly twice the global average (≈ 250 per 100,000)³. There has been an increase in the incidence of CHD, hypertension, and other cardiovascular conditions. Reports from several regional cardiology centers indicate a 10–15% rise in primary incidence rates over the past 5 years [3]. Concurrently, there has been an increase in the prevalence of risk factors (RFs) such as obesity, type 2 diabetes melli-

tus (T2DM), and metabolic syndrome, further exacerbating the epidemiological situation⁴. Recently, researchers have increasingly focused on comorbidity of somatic diseases as a key factor affecting work capacity and prognosis in patients with cardiovascular pathology [4]. This encompasses not only pathogenetically linked disorders but also independent diseases of various organs and systems, which adversely affect the course of CVDs [5].

Metabolic dysfunction-associated steatotic liver disease (MASLD) is a global health problem affecting up to 25% of the world's adult population [6]. Among patients with obesity and T2DM, the prevalence of MASLD reaches 70–90% [6]. Research indicates that MASLD is associated with an increased risk of cardiovascular events such as myocardial infarction (MI) and stroke, independent of conventional risk factors [7, 8]. According to the European Association for the Study of the Liver, MASLD is recognized not only as a liver disease but also as a significant cardiometabolic condition affecting the prognosis of patients with CHD [9, 10].

Contemporary studies demonstrate that MASLD increases the risk of atherosclerosis development and CHD progression through mechanisms involving systemic inflammation, oxidative stress, insulin resistance, and lipid metabolism disorders [11]. A large meta-analysis conducted by the Department of Medicine, University of Verona (Italy) revealed that patients with

¹ World Health Organization. Cardiovascular Diseases (CVDs). Fact Sheet, 2021. <https://www.who.int/news-room/fact-sheets/detail/cardiovascular-diseases-cvds>

² Agency of Statistics under the President of the Republic of Uzbekistan. Key Statistical Indicators of Healthcare in the Republic of Uzbekistan. <https://stat.uz/en/official-statistics> («Healthcare and social protection»)

³ World Health Organization (WHO). Noncommunicable Diseases (NCD) Country Profiles: Uzbekistan. Geneva: WHO; 2022. <https://www.who.int/teams/noncommunicable-diseases/surveillance/data/profiles-ncd>

⁴ International Diabetes Federation (IDF). IDF Diabetes Atlas. 10th ed. Brussels: 2, IDF; 2021. https://diabetesatlas.org/idfawp/resource-files/2021/07/IDF_Atlas_10th_Edition_2021.pdf

MASLD have a significantly higher risk of cardiovascular complications (CVCs), including atherosclerotic vascular lesions, compared to those without this condition [6]. Other researchers reached similar conclusions, notably the group led by S. Francque from the Department of Gastroenterology and Hepatology at the University of Antwerp (Belgium). They identified a strong association between the severity of liver fibrosis in MASLD and increased cardiovascular mortality [11]. Pathogenetically, this is attributed to systemic manifestations of metabolic dysfunction: heightened inflammatory response, activation of oxidative stress, development of insulin resistance, and progressive alterations in lipid metabolism, which collectively promote the formation and progression of atherosclerosis and CHD [4].

Asymmetric dimethylarginine (ADMA), an endogenous inhibitor of nitric oxide synthesis, plays a key role in the development of endothelial dysfunction, which underlies atherosclerosis and CHD [12]. ADMA levels increase in the presence of MASLD, as confirmed by data from large-scale studies (NHANES and ATHEROGEN) [13]. In patients with coexisting MASLD and CHD, ADMA levels may serve as a marker of cardiovascular risk [14]. However, there is insufficient data in the scientific literature regarding its clinical significance in this patient population.

Given the global epidemic of obesity and metabolic disorders, studying the impact of MASLD on CHD progression and related biochemical markers such as ADMA is of critical importance [15]. This will enable the improvement of diagnostic, therapeutic, and preventive strategies for CVCs in this patient group [16].

The aim of the study was to compare ADMA levels in patients with CHD with and without MASLD and to assess the potential association between ADMA levels and metabolic disorders.

Methods

A pilot, open-label clinical study was conducted at the Department of Cardiometabolic Disorders of the Republican Specialized Scientific and Practical Medical Center of Cardiology from July to December 2024. The study included 50 patients with CHD, equally divided into two groups: 25 patients with MASLD and 25 without MASLD.

Inclusion Criteria:

- Confirmed diagnosis of CHD (stable exertional angina, functional class I–II)

- Age 34–70 years
- Signed informed consent

Exclusion Criteria:

- Unstable angina, acute coronary syndrome, or history of myocardial infarction.
- Chronic heart failure NYHA class III–IV or left ventricular ejection fraction <40 %.
- Severe comorbidities affecting ADMA levels or endothelial function, such as chronic kidney disease (GFR <45 mL/min/1.73m²), liver cirrhosis or severe chronic liver disease, active malignancies, significant systemic inflammatory or autoimmune disorders.
- Decompensated type 1 diabetes, uncontrolled type 2 diabetes (HbA1c >9 % at enrollment, patients required treatment adjustment prior to assessment)
- Use of ADMA-affecting medications within 4 weeks prior to enrollment (e.g., high-dose L-arginine, PDE-5 inhibitors)

CHD was confirmed through clinical evaluation, electrocardiography (ECG), and coronary angiography (CAG) [2]. MASLD was diagnosed via liver ultrasound, supplemented by elastography for fibrosis staging when indicated. All participants underwent standardized laboratory testing, including complete blood count, comprehensive biochemistry panel, lipid profile, fasting glucose and insulin with HOMA-IR calculation. Serum ADMA levels were quantified using enzyme-linked immunosorbent assay (ELISA) with certified commercial kits (Beijing Solarbio Science & Technology Co., Ltd.).

Statistical analysis

Statistical analysis was performed using IBM SPSS Statistics software. Normally distributed data are presented as mean ± standard deviation, with between-group comparisons made using Student's t-test. For non-normally distributed data, the Mann-Whitney U test was applied. Statistical significance was defined as $p < 0.05$.

The study was conducted in accordance with the principles of the Declaration of Helsinki and approved by the local ethics committee. All patients received detailed information regarding study aims and potential risks, and signed an informed consent for participation and processing of their medical data.

Results

The study included 50 patients with CHD, divided into two groups: 25 with MASLD and 25 without MASLD.




Fig. 1. Distribution of patients by angina functional class

The mean age was 54.2 years in the MASLD group and 53.8 years in the non-MASLD group, with no statistically significant difference ($p = 0.73$). Males comprised 56 % of the MASLD group and 60 % of the non-MASLD group ($p = 0.64$). All patients had stable CHD classified as Canadian Cardiovascular Society (CCS) functional class (FC) I–II (Fig. 1). CHD diagnosis was established based on medical history, ECG, and CAG when clinically indicated. Patients received guideline-directed medical therapy including antianginal agents, antiplatelet therapy, and lipid-lowering treatment. Thus, group allocation was based exclusively on MASLD status, with comparable age-sex distributions and identical CHD functional class severity between groups.

Patients with MASLD had a significantly higher mean body mass index (BMI) (32.8 kg/m^2) compared to the non-MASLD group (27.1 kg/m^2 , $p < 0.01$). Arterial hypertension (AH) was present in 80 % of MASLD patients versus 48 % in non-MASLD patients ($p < 0.05$). T2DM was diagnosed in 52 % of MASLD patients and 20 % of non-MASLD patients ($p < 0.05$). Low-density lipoprotein (LDL) levels were higher in the MASLD

group (4.7 mmol/L) than in the non-MASLD group (3.1 mmol/L , $p < 0.01$) (Table 1).

Table 1. Clinical and Metabolic Characteristics of CHD Patients Stratified by MASLD Status

Parameter	CHD+MASLD (n=25)	CHD (n=25)	P
Mean age (years)	54.2 ± 6.1	53.8 ± 5.9	-
Male	14 (56 %)	16 (60 %)	-
BMI (kg/m^2)	32.8 ± 4.2	27.1 ± 3.5	<0.01
AH (%)	80 %	48 %	<0.05
T2DM (%)	52 %	20 %	<0.05
LDL (mmol/L)	4.7 ± 0.8	3.1 ± 0.7	<0.01

ADMA levels were significantly higher in MASLD patients compared to non-MASLD patients. The mean ADMA level in the MASLD group was $165.85 \pm 24.27 \text{ ng/mL}$, whereas the non-MASLD group showed significantly lower levels ($129.44 \pm 20.22 \text{ ng/mL}$). This difference was statistically significant ($p < 0.001$), indicating significant endothelial dysfunction in MASLD patients (Fig. 2).

Additional analysis revealed significantly higher ADMA levels in patients with a greater number of




Fig. 2. ADMA levels in study groups

metabolic disorders. Patients with 1–2 metabolic criteria (e.g., hypertension and obesity) had ADMA levels of 140 ± 15 ng/mL. Those with ≥ 3 criteria (e.g., obesity, type 2 diabetes, hypertension, dyslipidemia) showed substantially higher ADMA levels, averaging 190 ± 20 ng/mL ($p < 0.01$) (Table 2).

Table 2. Analysis of ADMA Levels According to Number of Metabolic Criteria

Study group	ADMA level (ng/mL)	P
1–2 metabolic criteria	140 ± 15	<0.01
3 and more criteria	190 ± 20	-

Discussion

This study presents the first domestic cohort analysis comparing ADMA levels in stable CHD patients stratified by MASLD status. Our findings confirm that MASLD comorbidity is associated with a pronounced elevation of this endothelial dysfunction biomarker. The observed increase (~ 36 ng/mL, $+28\%$) aligns with international studies [12]. For instance, Younossi et al. [7] reported 0.09 $\mu\text{mol/L}$ higher ADMA concentrations (~ 23 – 25 ng/mL) in biopsy-confirmed MASLD patients versus controls. Recent MASLD reviews indicate a mean excess of 0.1 $\mu\text{mol/L}$. Thus, our data demonstrate consistent manifestation of this phenomenon across diverse ethnic and geographic populations.

The causal relationship between MASLD and elevated ADMA is largely mediated by insulin resistance and chronic systemic inflammation [11]. Insulin resistance suppresses dimethylarginine dimethylaminohydrolase (DDAH) activity, the enzyme responsible for ADMA degradation, leading to its accumulation. Concurrently, proinflammatory cytokines (TNF- α , IL-6) and oxidative stress in the liver and vascular wall further inhibit DDAH while enhancing arginine methylation in proteins, thereby increasing ADMA synthesis [5].

ADMA itself competitively inhibits endothelial nitric oxide synthase, reducing nitric oxide bioavailability, enhancing vascular tone, and promoting atherogenesis. This mechanism explains the more frequent and severe coronary artery involvement in patients with combined MASLD and CHD [12].

Notably, in our cohort, ADMA levels exhibited a proportional increase with the number of metabolic dis-

orders: patients with ≥ 3 criteria showed 35% higher values than those with 1–2 criteria. A similar gradient was demonstrated in untreated hypertensive patients (ADMA $\uparrow \sim 20\%$), reinforcing the concept of an ADMA-NO “metabolic cascade”. Consequently, ADMA may serve as an integrative biomarker of cumulative metabolic burden on the endothelium [14]. The elevated ADMA levels observed in MASLD patients have significant clinical relevance: they enable more accurate cardiovascular risk stratification in stable CHD patients where conventional scores (SCORE2, GRACE) underestimate hepatic comorbidity impact [7]; underscore the necessity for early lifestyle modification and pharmacological management of insulin resistance to reduce ADMA and improve vascular prognosis; and function as a treatment response biomarker, where ADMA reduction during statin or combination lipid-lowering therapy correlates with atherosclerotic plaque regression [1].

As a pilot investigation, this study has inherent limitations: its single-center design and restricted sample size reduce the generalizability of findings. The single-timepoint measurement of ADMA precludes assessment of its dynamics in response to therapeutic interventions. The absence of instrumental endothelial function evaluation (flow-mediated dilation, FMD) [10] limits direct validation of ADMA as a surrogate marker. Future large-scale prospective studies employing multivariate models (incorporating ADMA + FMD + liver fibrosis indices) are needed to determine whether including ADMA enhances predictive accuracy for CVCs in MASLD patients. Furthermore, therapeutic strategies targeting ADMA reduction, such as pharmacologic DDAH activation or L-citrulline supplementation [15], remain promising yet underexplored avenues in cardiometabolic pharmacology.

Conclusion

This study demonstrated significantly higher ADMA levels in CHD patients with comorbid MASLD compared to those without MASLD. A positive correlation was observed between the number of metabolic disorders (obesity, hypertension, type 2 diabetes) and elevated ADMA concentrations, suggesting its potential role in the pathogenesis of endothelial dysfunction and CVCs. These findings support the need for a personalized approach to diagnosing and treating CHD patients that accounts for concomitant meta-

- 16 Nurillaeva N.M., Makhkamova M.M., Srojidinova N.Z.
The role of metabolic disorders in the formation of asymmetric dimethylarginine levels in patients...
DOI: 10.24412/2311-1623-2025-47-13-20
-

bolic disorders, while highlighting the potential utility of ADMA as a cardiovascular risk stratification biomarker.

Conflict of interest: none declared.

References

- Lewalter T, Becher PM, Bauersachs J, et al. Therapeutic Consequences and Prognostic Impact of Multimorbidity in Real-World Heart-Failure Patients. *J Clin Med*. 2024;14(1):p139. DOI:10.3390/jcm14010139
- European Society of Cardiology (ESC). ESC Guidelines on Cardiovascular Disease Prevention in Clinical Practice. *Eur. Heart J*. 2021, 42, 3227–3337. DOI: 10.1093/eurheartj/ehab484
- Mamedov MN, Deev AD, Mekhdiyev SKh Priorities of primary prevention of cardiovascular disease: the results of multicenter international cohort study AHS I (Azerbaijan Heart Study, part I). *International Journal of Heart and Vascular Diseases*. 2018. 6(18): 3–11. Russian.
- Makhkamova MM, Nurillayeva NM Relationship between metabolic associated steatotic liver disease and ischemic heart disease: clinical aspects of comorbidity. *Cardiology journal of Uzbekistan* 2024. 30;1(4): 221–7.
- Dowsett L, Higgins E, Alanazi S et al. ADMA: A Key Player in the Relationship between Vascular Dysfunction and Inflammation in Atherosclerosis. *Journal of clinical medicine*. 2020. 9(9): 3026. DOI: 10.3390/jcm9093026
- Targher G, Byrne CD, Tilg H. NAFLD and increased risk of cardiovascular disease: clinical associations, pathophysiological mechanisms, and pharmacological implications. *Gut*. 2020;69(9): 1691–1705. DOI: 10.1136/gutjnl-2020-320622
- Younossi Z, Anstee QM, Marietti M, et al. Global burden of NAFLD and NASH: trends, predictions, risk factors and prevention. *Nat Rev Gastroenterol Hepatol*. 2018;15(1):11–20. DOI: 10.1038/nrgastro.2017.109
- Medvedev VM, Tikhonova AS, Solovyov AV. Metabolic Dysfunction in Ischemic Heart Disease: Association with Inflammation and Lipid Metabolism. *Medical News*. 2022; 31(8): 45–51.
- Pauly DF, Pepine CJ Ischemic heart disease: metabolic approaches to management. *Clinical cardiology*. 2024; 27(8): 439–441. DOI: 10.1002/clc.4960270802
- European Association for the Study of the Liver (EASL), European Association for the Study of Diabetes (EASD), European Association for the Study of Obesity (EASO). EASL–EASD–EASO Clinical Practice Guidelines for the management of non-alcoholic fatty liver disease. *J Hepatol*. 2019;64(6):1388–1402. DOI: 10.1016/j.jhep.2015.11.004
- Francque S, van der Graaff D, Kwanten W. Non-alcoholic fatty liver disease and cardiovascular risk: Pathophysiological mechanisms and implications. *Journal of Hepatology*. 2019; 70(4): 749–767. DOI: 10.1016/j.jhep.2018.12.022
- Rodionov RN, Blokhin IO, Galagudza MM et al. Asymmetric dimethylarginine and its role in the etiology and pathogenesis of cardiovascular diseases. *Arterial Hypertension*. 2019; 14(4): 306–314. DOI: 10.1161/01.ATV.0000081742.92006.59
- Baum C, Johannsen SS, Zeller T et al. Gutenberg Health Study investigators ADMA and arginine derivatives in relation to non-invasive vascular function in the general population. *Atherosclerosis*. 2016; 244: 149–156. DOI: 10.1016/j.atherosclerosis.2015.10.101
- Zhu W, Huang X, Li C et al. Asymmetric Dimethylarginine and All-Cause Mortality: A Systematic Review and Meta-Analysis. *Sci. Rep*. 2020; 10: 8672. DOI: 10.1038/s41598-020-65125-y
- Guo X, Yin X, Liu Z, Wang J. Non-Alcoholic Fatty Liver Disease (NAFLD) Pathogenesis and Natural Products for Prevention and Treatment. *Int. J. Mol. Sci*. 2022; 23: 15489. DOI: 10.3390/ijms232415489
- Klaritskaya IL, Bondar KG Metabolically associated fatty liver disease and comorbid conditions. *Crimean Therapeutic Journal*. 2024; 4: 13–27. Russian.

Prognostic value of selected biomarkers in comorbid patients with chronic heart failure after COVID-19

Kovalenko E.V., Belaya O.L., Markova L.I., Evdokimov V.V.

Federal Scientific State Budgetary Institution "N.A. Semashko National Research Institute of Public Health",
Moscow, Russia.

AUTHORS

Elena V. Kovalenko*, MD, PhD, Professor of the Department of Hospital Therapy № 2, Federal Scientific State Budgetary Institution "N.A. Semashko National Research Institute of Public Health", Moscow, Russia. ORCID: 0000-0001-9202-3522

Olga L. Belaya, MD, PhD, Professor of the Department of Hospital Therapy № 2, Federal Scientific State Budgetary Institution "N.A. Semashko National Research Institute of Public Health", Moscow, Russia. ORCID: 0000-0002-5256-3580

Lyudmila I. Markova, MD, PhD, Professor of the Department of Hospital Therapy № 2, Federal Scientific State Budgetary Institution "N.A. Semashko National Research Institute of Public Health", Moscow, Russia. ORCID: 0000-0002-3396-9235

Vladimir V. Evdokimov, MD, PhD, Professor, Head of the Department of Hospital Therapy № 2, Federal Scientific State Budgetary Institution "N.A. Semashko National Research Institute of Public Health", Moscow, Russia. ORCID: 0000-0003-3910-2488

The aim of the study was to evaluate the role of such biomarkers as N-terminal prohormone of brain natriuretic peptide (NT-proBNP), cystatin C, and galectin-3 for prediction of poor outcomes of chronic heart failure with preserved ejection fraction (HFpEF) of the left ventricle (LV) in patients with type 2 diabetes mellitus (T2DM) and chronic kidney disease (CKD) with a history of COVID-19.

Methods. This study included 135 patients with HFpEF and cardiovascular disorders, T2DM, CKD, and a history of COVID-19 from more than 3 months prior to enrollment. Patients were assessed clinically, instrumentally and

using laboratory methods, including NT-proBNP, cystatin C, and galectin-3 assays and echocardiography. The occurrence of heart failure (HF) decompensation and cardiovascular death (CVD) was recorded over 18 months.

Results. A multivariate regression analysis was used to build a basic model for estimation of probability of HF decompensation and CVD in patients with a history of COVID-19, HFpEF, T2DM, and CKD and included risk factors, such as rating scale of clinical state (ShOKS), stroke volume, left ventricular mass index, E/e' ratio, and NT-proBNP, which showed a high prognostic potential

* Corresponding author. Tel. +7 (915) 262-89-82. E-mail: elkovalenko76@mail.ru

(AUC=0.856; 95% CI: 0.717–0.994; $p<0.001$, sensitivity 85.6%, specificity 79.2%). Adding cystatin C and galectin-3 improved prognostic properties of the resulting models (AUC=0.867; 95% CI: 0.808–0.923, $p<0.001$, sensitivity 86.3%, specificity 75% and AUC=0.866; 95% CI: 0.809–0.923, $p<0.001$, sensitivity 84.9%, specificity 83.3%, respectively). The best model was developed using levels of assessed biomarkers (AUC=0.879; 95% CI: 0.845–0.913, $p<0.001$, sensitivity 87.5%, specificity 88.8%).

Conclusion. Including NT-proBNP, cystatin C, and galectin-3 into a multifactorial model can improve the prediction of HF worsening and CVD in patients with a history of COVID-19, HFpEF, T2DM, and CKD.

Keywords: chronic heart failure, NT-proBNP, galectin-3, cystatin C, COVID-19, prognosis.

Conflict of interest: none declared.

Accepted: 24.06.2025

Received: 04.08.2025



For citation: Kovalenko E.V., Belaya O.L., Markova L.I., Evdokimov V.V. Prognostic value of selected biomarkers in comorbid patients with chronic heart failure after COVID-19. *International Journal of Heart and Vascular Diseases*. 2025. 13(47):17-25. DOI: 10.24412/2311-1623-2025-47-21-32

Introduction

Many chronic heart failure (CHF) patients have a preserved ejection fraction (EF) of the left ventricle (LV). These patients often have arterial hypertension (AH), type 2 diabetes mellitus (T2DM), obesity, coronary heart disease (CHD), chronic kidney disease (CKD), and are elderly [1]. These comorbidities mutually exacerbate each other, regardless of their order of appearance [2].

Long-lasting impact of established risk factors (RF) in comorbid patients lead to systemic inflammation, atherosclerosis, and ischemia. Against the background of metabolic and hemodynamic disorders, associated with insulin resistance, renin angiotensin aldosterone system (RAAS) activation, elevated levels of glycation end products and adipokines, patients experience oxidative injury of cardiomyocytes, glomerular filter structures, tubular epitheliocytes, mesangial cells, accompanied by increased active transforming growth factor β production, stimulation of collagen formation and fibrosis. Structural changes in heart and kidneys result in pathologic remodeling and dysfunction. In turn, cardiorenal dysfunction leads to compensatory activation of RAAS and establishes a vicious circle [3]. Thereby, inflammation and fibrosis compose the basis of cardiorenal dysfunction formation and progression pathogenesis in patients with CHF, T2DM, and CKD. Therefore, biomarkers reflecting these processes hold promise for prognostic assessment and represent a sought-after area of research/medicine.

Cystatin C is one of well-studied markers in cardiovascular and renal disorder, which is a member of cysteine proteinase inhibitors family. Serum cystatin C levels are known to represent glomerular filtration rate (GFR) more accurately, than serum creatinine. In addition to renal dysfunction, high cystatin C concentration might be associated with inflammatory and atherogenic processes [4]. Levels of this marker were studied during the coronavirus infection pandemic (COVID-19). The meta-analysis revealed an association between high serum concentrations of this biomarker and both COVID-19 infection severity and mortality in hospitalized patients [5]. Nevertheless, the significance of cystatin C in patients after the COVID-19 infection is yet to be defined.

In the past few years, the possibility of the use of galectin-3 in patients with cardiovascular disorder, including CHF. Controversial study results preclude its recommendation as a reliable indicator of heart failure (HF). However, the use of galectin-3 in combination with gold-standard HF biomarkers (B-type natriuretic peptide (BNP)/N-terminal pro-BNP (NT-proBNP) might improve the prognostic assessment of adverse events [6]. The significance of galectin-3 in the pathogenesis of cardiovascular remodeling, inflammatory and autoimmune processes, and neoplasia [7]. Moreover, the increase in galectin-3-positive macrophages in renal glomeruli in T2DM is associated with level of albuminuria, progression of kidney dysfunction, and a poor prognosis [8]. For the time

being, galectin-3 is considered one of the markers of fibrosis and inflammation [9].

Most studies on multiple-marker prognostic assessment were conducted before the COVID-19 pandemic, and its delayed consequences are still to be researched. It has been established that patients with CHD, AH, T2DM, CKD, and CHF fall into the category of poor COVID-19 outcomes and very high risk of cardiovascular complications [10, 11]. Despite the significant evidence base of clinical application of various biomarkers, the question of the gold-standard indicator for long-term CHF poor prognosis in patients with cardiorenal dysfunction after COVID-19 remains open [12].

Aim of the study

The aim of the study is to estimate the role of biomarkers, such as NT-proBNP, cystatin C, and galectin-3 for prognostic assessment of poor outcomes in patients with HFpEF, T2DM, CKD, and COVID-19 in medical history.

Methods

The study included 135 stable patients with HFpEF, T2DM, CKD, and a confirmed COVID-19 from 3 or more months prior to the hospitalization. All participants signed an informed consent form. The median age was 70 [62, 74]. The study included patients with CHF of I–III NYHA functional class with medical history of grade 1–2 of AH, and 85 (76.58%) participants had CHD. The exclusion criteria were acute myocardial infarction, stroke or a transient ischemic attack for 6 months prior to the enrollment, IV NYHA functional class of CHF, hemodynamically significant cardiac abnormalities, grade 3 AH, severe renal dysfunction with estimated glomerular filtration rate (eGFR) < 25 mL/min/1.73 m² by the Chronic Kidney Disease Epidemiology Collaboration Formula (CKD-EPI) 2012, level of glycated hemoglobin (HbA1c) > 12%. The period of observation was 18 months. NYHA functional class of CHF, clinical severity assessed by the Rating Scale of Clinical State, Kaplan-Feinstein comorbidity index, 6-minute walk test distance (6MWT), echocardiographic data (EchoCG), eGFR by the CKD-EPI 2012, urine albumin-to-creatinine ratio (uACR), serum levels of NT-proBNP, galectin-3, and cystatin C assessment. Transthoracic EchoCG with myocardial tissue Doppler imaging was carried out. Left ventricular ejection fraction was estimated

with Simpson's method of disks. Diastolic function of the left ventricle was estimated using transmitral flow in pulsed-wave and tissue Doppler modes. Biomarker levels were quantitatively assessed with Enzyme-Linked Immunosorbent Assay (ELISA) with the following kits: "Biomedica NT-proBNP" (Austria), "human Galectin-3 ELISA", Bender MedSystems (Austria), "Human Cystatin C ELISA", BioVendor (Czech Republic). Additionally, a group of 21 participants (12 women and 9 men) with no signs of disease and no medical history of COVID-19 for to establish control biomarker levels. This group had a median age of 54 [49, 58] and median levels of NT-proBNP 77 [49, 91] pg/mL, galectin-3 – 2.7 [1.8, 3.6] ng/mL, cystatin C 0.781 [0.694, 0.820] mg/L.

The Inter-University Committee on ethics from 20.01.2022 (protocol № 01–22) approved the study.

Statistical analysis

Statistical data were processed using Python programming language (3.10 version, scipy, statmodels, pandas, math libraries) and MedCalc software. Descriptive statistics for group parameters included frequencies and percentages for categorical variables, and medians with interquartile ranges (Q25, Q75) for continuous variables, accounting for the distribution type. The Kolmogorov-Smirnov method was used for normality tests. Differences in proportions of a trait between groups were assessed using the proportion Z-test. Differences in medians were evaluated with the Kruskal-Wallis test. The strength of association between continuous variables was measured using Spearman's correlation, while differences between categorical variables was assessed with Fisher's exact test. The association between dichotomous and continuous variables was examined using point-biserial correlation. Risk was estimated by calculating the odds ratio and relative risk. Differences in indicators between groups were considered statistically significant at $p < 0.05$. Cut-off values for the concentrations of the investigated markers, along with their sensitivity and specificity, were derived from ROC analysis. A predictive model for adverse events of the composite endpoint (CEP), including cases of cardiovascular death (CVD) and HF decompensation, was created using multivariate binary logistic regression analysis. The sensitivity and specificity of the predictive model were determined via ROC analysis.

Results

According to medical history, the median duration of CHF, AH, T2DM, and CKD was 5.0 [3.0, 8.0] years, 19 [17, 21] years, 12 [9, 17] years, and 2 [2, 3] years, respectively. Among the 85 (76.58%) patients with CHD, 49 (36.3%) had two or more affected coronary arteries, 34 (30.63%) had a history of myocardial infarction, 38 (34.23%) had undergone percutaneous coronary intervention, and 7 (6.31%) had undergone coronary artery bypass grafting. Preproliferative and proliferative stages of diabetic retinopathy were identified in 42 (31.11%) and 10 (7.41%) patients, respectively. Distal diabetic neuropathy was diagnosed in 126 (93.33%) patients, of whom 54 (40.0%) had sensorimotor neuropathy. The median Kaplan-Feinstein comorbidity index score was 15 [12, 16] points. The majority of included patients, 87 (66.44%), had NYHA FC II of CHF, 46 (34.07%) patients had NYHA FC III of CHF, and 2 (1.48%) had NYHA FC I of CHF. Key clinical, functional, and laboratory parameters are presented in Table 1.

Table 1. Clinical, functional, and laboratory parameters, median [Q25, Q75]

Parameter	Value
BMI, kg/m ²	31.51 [29.7, 34.41]
Waist circumference, cm	105.0 [98.5, 115.0]
ShOKS (Rating Scale of Clinical State, Russian), points	5.0 [4.0, 7.0]
6MWT (6-Minute Walk Test), m	348.0 [285.0, 395.0]
HbA1c, %	7.8 [7.1, 9.1]
Cholesterol, mmol/L	4.45 [3.78, 4.99]
Sodium, mmol/L	140.2 [139.3, 141.6]
Potassium, mmol/L	4.53 [3.91, 5.04]
Urea, mmol/L	7.17 [6.1, 9.45]
Creatinine, μ mol/L	105.0 [92.0, 123.9]
Creatinine eGFR, mL/min/1.73 m ²	51.89 [39.67, 69.39]
Cystatin C, mg/L	1.54 [1.26, 2.08]
Cystatin eGFR, mL/min/1.73 m ²	39.0 [27.49, 53.03]
uACR, mg/g	150.0 [66.0, 297.0]
NT-proBNP, pg/mL	679.0 [452.0, 911.0]
Galectin-3, ng/mL	11.52 [9.4, 14.58]

All patients had concentrations of NT-proBNP, galectin-3, and cystatin C above control values. An analysis of the levels of the studied biomarkers was performed across different ranges of creatinine-based eGFR, corresponding to CKD stages 1 through 4 (Figures 1–3).




Fig. 1. NT-proBNP levels depending on eGFR, $p=0.123$





Fig. 2. Galectin-3 levels depending on eGFR, $p<0.001$



Unlike NT-proBNP concentration, showing no significant difference across the subgroups ($p=0.123$), galectin-3 and cystatin C levels increased significantly with worsening renal dysfunction ($p<0.001$).

Correlation analysis revealed moderate positive correlations between cystatin C level and CHF duration ($r=0.325$, $p=0.0001$), Kaplan-Feinstein comorbidity index ($r=0.449$, $p<0.0001$), Left ventricular myocardial mass index (LVMMI) ($r=0.442$, $p<0.0001$), NT-proBNP concentration ($r=0.498$, $p<0.0001$). Strong positive correlations were found between cystatin C and Creatinine levels ($r=0.717$, $p<0.0001$), Galectin-3 levels ($r=0.777$, $p<0.0001$). Significant negative correlations existed between cystatin C and 6MWT distance ($r=-0.456$, $p<0.0001$) and LVEF ($r=-0.430$, $p<0.0001$).

Galectin-3 concentration correlated with T2DM duration ($r=0.371$, $p<0.0001$), HF duration ($r=0.323$, $p=0.00013$), Kaplan-Feinstein comorbidity index ($r=0.429$, $p<0.0001$), 6MWT distance ($r=-0.417$, $p<0.0001$), uACR level ($r=0.345$, $p<0.0001$), Creatinine level ($r=0.603$, $p<0.0001$), eGFR ($r=-0.490$, $p<0.0001$), LVEF ($r=-0.363$, $p<0.0001$), LVMMI ($r=0.361$, $p<0.0001$), E/e' ratio ($r=0.519$, $p<0.0001$), NT-proBNP level ($r=0.533$, $p<0.0001$). As the analysis reveals, markers reflecting diverse processes within the complex pathogenesis of CHF in comorbid patients, and parameters of functional status, target organ damage severity, and duration of risk factor exposure, are closely associated.

Using point-biserial correlation, a significant association was established between the severity of prior COVID-19 and levels of the studied biomarkers: NT-proBNP ($r=0.238$, $p=0.005$), galectin-3 ($r=0.371$, $p<0.0001$), and cystatin C ($r=0.279$, $p=0.001$). The stronger correlation of prior COVID-19 severity with galectin-3 concentration may be attributed to patients' initially more severe clinical status or consequences of the infectious process.

During the period of observation, CEP events occurred in 54 patients (40%): 3 (2.2%) patients died from cardiovascular causes, 23 (17.03%) patients were hospitalized due to CHF decompensation, 28 (20.74%) patients required outpatient medication adjustment for worsening HF symptoms. Assessment of CEP cases revealed that adverse outcomes occurred more frequently in patients with higher baseline func-

tional impairment, higher Kaplan-Feinstein comorbidity index (16.0 points [15.0, 18.0] compared to 14.0 points [11.0, 15.0] in patients without CEP, $p<0.001$; Table 2), more pronounced cardiorenal dysfunction. Moreover, 41 patients (75.93%) who experienced CEP had a CHF hospitalization within the 12 months prior to the study, compared to 28 patients (34.57%) in the non-CEP subgroup, $p<0.001$.

Table 2. Baseline characteristics of patients stratified by composite endpoint occurrence

Parameter	No CEP (n=81)	CEP (n=54)	p-value
Age, years	70 [61; 73]	71 [65; 75]	0.16
Male, n [%]	33 [40.74]	31 [57.41]	0.04
CHF duration, years	4 [2; 6]	7 [5; 12]	<0.001
T2DM duration, years	10 [9; 17]	14 [11; 19]	0.002
CKD duration, years	2 [1; 3]	3 [2; 4]	<0.001
BMI, kg/m ²	31.38 [29.94; 34.09]	31.51 [29.07; 35.02]	0.59
ShOKS, points	5 [4; 6]	7 [6; 8]	<0.001
6MWT, m	380 [325; 405]	290 [269; 365]	<0.001
Creatinine, µmol	98.0 [82.5; 122.0]	108.7 [96.0; 134.0]	0.003
uACR, mg/g	122.3 [66.0; 263.0]	263.0 [75.0; 526.0]	0.007
NT-proBNP, pg/mL	512 [410; 746]	897 [690; 1112.0]	<0.001
galectin-3, ng/mL	10.5 [8.74; 12.3]	14.94 [11.4; 17.01]	<0.001
Cystatin C, mg/L	1.4 [1.16; 1.79]	2.02 [1.53; 2.41]	<0.001
Comorbidities, n (%)			
CHD	57 [70.37]		
Myocardial infarction in medical history	16 [19.75]	46 [85.19]	0.04
Atrial fibrillation	21 [25.93]	25 [46.3]	0.001
Obstructive sleep apnea	3 [3.7]	29 [53.7]	0.001
		7 [12.96]	0.04
LVEF, %	54 [52; 57]	53 [51; 56]	0.03
Left atrial volume index, mL/m ²	34.67 [34.17; 35.6]	36.96 [35.1; 45.26]	<0.001
LVMMI, g/m ²	139.25 [116.96; 153.45]	151.03 [132.53; 164.19]	0.001
E/e'	14.18 [10.6; 15.6]	16.48 [15.26; 17.79]	<0.001

When assessing the prognostic significance of NT-proBNP, cystatin C, and galectin-3, the markers were tested in ROC analysis. The area under the

Table 3. Prognostic characteristics of investigated markers in univariate analysis

Parameter	AUC	95%CI	Threshold	Sensitivity, %	Specificity, %	p
NT-proBNP, pg/mL	0.787	0.633–0.942	897.0	82.5	73	<0.001
Cystatin C, mg/L	0.788	0.634–0.945	1.683	83.3	73.6	<0.001
galectin-3, ng/mL	0.791	0.637–0.949	13.08	84.1	72.7	<0.001

curve (AUC) was 0.787, 0.788, and 0.791, respectively. Marker levels above established cut-off values – NT-proBNP: 897 pg/mL [relative risk (RR) = 4.523, 95 % confidence interval (CI): 3.277–5.77; $p<0.001$], cystatin C: 1.683 mg/L (RR=6.428, 95 % CI: 5.735–7.122; $p<0.001$), galectin-3: 13.08 ng/mL (RR=3.214, 95 % CI: 2.103–4.325; $p<0.001$) identified patients as having an elevated CEP risk (Table 3).

Using multivariate logistic regression analysis, baseline mathematical Model 1 was constructed to predict the probability of HF decompensation and CVD, incorporating the ShOKS score, stroke volume (SV), LVMMI, E/e' ratio, and NT-proBNP concentration (Table 4). The resulting model demonstrated high prognostic value (AUC = 0.856; 95 % CI: 0.717–0.994; $p<0.001$; sensitivity 85.6 %, specificity 79.2 %) (Fig. 4).

Table 4. Risk factors for HF decompensation in multivariate regression models (Models 1, 2)

Parameter	Model 1 (AUC=0.856; $p<0.001$)		Model 2 (AUC=0.867; $p<0.001$)	
	OR (95%CI)	p-value	OR (95%CI)	p-value
NT-proBNP	1.01 (1.00–1.02)	0.001	1.01 (1.00–1.02)	<0.001
ShOKS	1.06 (0.77–1.46)	0.335	1.01 (0.72–1.41)	0.933
SV (Stroke volume)	0.98 (0.94–1.02)	0.929	0.99 (0.95–1.03)	0.73
LVMMI	0.96 (0.94–0.99)	0.045	0.96 (0.93–0.99)	0.01
E/e'	1.14 (0.97–1.34)	0.194	1.09 (0.91–1.29)	0.311
Cystatin C	No data	No data	2.52 (0.78–8.09)	0.199




Fig. 4. ROC curve for baseline Model 1 including NT-proBNP

When assessing the prognostic capability of the studied biomarkers, cystatin C and galectin-3 concentrations were sequentially added to the baseline model and the performance of the resulting models

was analyzed. Incorporating cystatin C into the original model (Model 2) improved the prognostic properties of Model 1, increasing the area under the ROC curve (AUC=0.867; 95 % CI: 0.808–0.923, $p<0.001$; sensitivity 86.3 %, specificity 75 %) (Fig. 5, Table 4).




Fig. 5. ROC curve for Model 2 including NT-proBNP and cystatin C

Model 3, which included galectin-3 concentration (AUC=0.866; 95 % CI: 0.809–0.923, $p<0.001$; sensitivity 84.9 %, specificity 83.3 %), demonstrated a larger AUC compared to baseline Model 1 but did not significantly differ in prognostic significance from the model containing cystatin C (Fig. 6, Table 5).




Fig. 6. ROC curve for Model 3 including NT-proBNP and galectin-3

The inclusion of all studied biomarkers further improved the predictive properties of the model (AUC=0.879; 95 % CI: 0.845–0.913, $p<0.001$; sensitivity 87.5 %, specificity 88.8 %) (Fig. 7, Table 5). Thus, compared to the baseline model incorporating NT-proBNP concentration, the addition of cystatin C

Table 5. Risk factors for HF decompensation in multivariate regression models (Models 3, 4)

Parameter	Model 3 (AUC=0.866; p<0.001)		Model 4 (AUC=0.879; p<0.001)	
	OR (95%CI)	p-value	OR (95%CI)	p-value
NT-proBNP	1.01 (1.00–1.02)	0.01	1.01 (1.0–1.02)	0.004
ShOKS	1.07 (0.77–1.49)	0.647	1.01 (0.7–1.43)	0.966
SV	0.98 (0.94–1.02)	0.516	0.99 (0.95–1.03)	0.74
LVMMI	0.96 (0.93–0.99)	0.016	0.96 (0.94–0.99)	0.021
E/e ⁺	1.08 (0.9–1.3)	0.363	0.99 (0.83–1.18)	0.965
Galectin-3	1.1 (0.92–1.31)	0.26	1.14 (0.92–1.4)	0.21
Cystatin C	No data	No data	1.47 (0.35–6.13)	0.592




Fig. 7. ROC curve for Model 4 including NT-proBNP, cystatin C, and galectin-3

and/or galectin-3 enhances the predictive capability for assessing the risk of HF decompensation and CVD in post-COVID-19 patients with HFpEF, T2DM, and CKD.

Discussion

The comprehensive algorithm for HF patients' diagnosis and treatment efficacy evaluation involves the use of biomarkers and EchoCG data. Extensive evidence has established NT-proBNP as a reference-standard biomarker in HF [13]. Study results confirmed the reasonably high prognostic value of NT-proBNP (ROC AUC=0.787; 95% CI 0.633–0.942, p<0.001) in HFpEF patients with cardiovascular disease, T2DM, CKD, and a history of COVID-19. The biomarker was a significant predictor in the multivariate model (p=0.001).

Given the scarcity of studies evaluating long-term prognosis using markers reflecting different aspects of HFpEF pathogenesis in patients with T2DM and CKD after COVID-19, this work investigated the prog-

nostic influence of established markers such as cystatin C and galectin-3 in comorbid HFpEF patients.

The study revealed correlations of varying strength between cystatin C levels and cardiac structural and functional parameters, indicating a close connection of cardiorenal dysfunction in HFpEF and CKD patients. The observed increase in cystatin C concentration corresponding to declining eGFR levels and severity of prior COVID-19 likely reflects more severe renal impairment and inflammation. In previous studies, cystatin C has proven to be an ideal marker for GFR, with its levels also associated with acute kidney injury and inflammatory status severity during the COVID-19 pandemic [4, 14, 15]. Elevated cystatin C in COVID-19 patients alongside reduced GFR may result from its involvement in the release of cytokines such as tumor necrosis factor- α (TNF- α), interleukin-12 (IL-12), interleukin-10 (IL-10), and nitric oxide (NO) [16]. Cystatin C activates the inducible isoform of nitric oxide synthase (iNOS), which generates higher NO concentrations compared to endothelial NOS (eNOS) and neuronal NOS (nNOS) during local and systemic inflammatory responses. This leads to the formation of highly reactive derivatives that cause irreversible alterations in intracellular structures and cellular apoptosis. This process represents one of the pathophysiological mechanisms underlying cytokine storm development and severe COVID-19 progression [17–19].

In prognostic assessment, cystatin C demonstrated predictive capability for the CEP development comparable to NT-proBNP. The established cystatin C cut-off value of ≥ 1.683 mg/L in the univariable model identified high-risk individuals for HF decompensation and CVD among clinically stable HFpEF patients with T2DM and CKD after COVID-19. Adding this biomarker to baseline parameters, which included NT-proBNP, improved prediction accuracy. Previous studies associated cystatin C levels >0.80 – 1.6 mg/L with increased mortality risk [20–22]. The higher cut-off value observed in this study may reflect the more severe clinical profile of our patient cohort.

In this study, galectin-3, being a biomarker of inflammation and fibrosis, exhibited correlations with T2DM duration, HF duration, patient comorbidity burden and functional status, left ventricular remodeling and diastolic dysfunction parameters, NT-proBNP levels, renal dysfunction markers (ACR, creatinine, and eGFR). Prolonged systemic exposure to risk fac-

tors evidently activates proinflammatory processes, oxidative stress, and fibrosis in target organs, leading to progressive functional decline. Similar but stronger correlations were reported in a Turkish study of HFpEF patients, where galectin-3 correlated with NT-proBNP ($r=0.90$, $p<0.0001$), left atrial volume index ($r=0.75$, $p=0.0001$), LVMMI ($r=0.86$, $p=0.0001$), E/e' ratio ($r=0.80$, $p=0.0001$) [23]. A weaker galectin-3/NT-proBNP correlation ($r=0.21$, $p=0.048$) was observed in another HFpEF cohort [24]. Association between galectin-3 and LV diastolic dysfunction was documented in HFpEF patients with ST-segment elevation myocardial infarction as well [25]. The observed link between galectin-3 and renal impairment severity aligns with prior research [8]. Impaired renal clearance, particularly in patients with kidney dysfunction, is presumed to contribute to elevated galectin-3 in HF [26].

In this study, a galectin-3 level ≥ 13.08 ng/mL (sensitivity 84.1%, specificity 72.7%, $p<0.001$) was associated with increased risk of the CEP. Previous studies also established statistically significant associations between galectin-3 and mortality in CHF patients [27]. Elevated galectin-3 levels were linked to worse outcomes regardless of treatment or NT-proBNP concentration [28]. Furthermore, in patients from CHF-risk groups, galectin-3 independently predicted HF development and 10-year mortality [29].

Model 4, which incorporated NT-proBNP, cystatin C, and galectin-3 in addition to baseline parameters, demonstrated the strongest predictive performance. Multimarker risk assessment for HF decompensation and CVD enhanced model performance, increasing

the ROC AUC to 0.879 ($p<0.001$). Thus, biomarkers reflecting core pathophysiological mechanisms of cardiorenal interactions in post-COVID-19 HFpEF patients with T2DM and CKD enable precise identification of high-risk individuals, facilitating timely therapeutic management for subclinical disease control and prevention of potential complications.

Conclusion

Analysis of this HFpEF cohort with T2DM and CKD demonstrated that measuring cystatin C and galectin-3 alongside NT-proBNP enhances risk stratification, addressing the urgent unmet need for more precise prediction tools for HF decompensation and CVD in comorbid post-COVID-19 patients. These findings reveal the potential clinical utility of inflammation and fibrosis biomarkers in comorbid HFpEF populations. Prospectively, serial monitoring of cystatin C and galectin-3, particularly in post-COVID patients, combined with multimarker assessment would provide refined prognostic insights and enable therapy efficacy evaluation. Extended follow-up of comorbid HFpEF patients using biomarkers reflecting diverse pathophysiological pathways, alongside developing specialized diagnostic and prognostic panels, may facilitate timely treatment correction. The stronger association between galectin-3 levels and COVID-19 severity warrants further investigation for objectively assessing post-infectious sequelae in critically ill patients.

Conflict of interest: none declared.

References

1. Polyakov DS, Fomin IV, Belenkov YuN, et al. Chronic heart failure in the Russian Federation: what has changed over 20 years of follow-up? Results of the EPOCH-CHF study. *Kardiologiya*. 2021; 61(4):4-14. Russian. 61(4):4-14. DOI: 10.18087/cardio.2021.4.n1628
2. Lawson CA, Testani JM, Mamas M et al. Chronic kidney disease, worsening renal function and outcomes in a heart failure community setting: a UK national study. *Int J Cardiol*. 2018; 267:120-127. DOI: 10.1016/j.ijcard.2018.04.090
3. Patel RN, Sharma A, Prasad A et al. Heart Failure With Preserved Ejection Fraction With CKD: A Narrative Review of a Multispecialty Disorder. *Kidney Med*. 2023; 5(12):100705. DOI: 10.1016/j.xkme.2023.100705
4. Belaya OL, Orangereeva VN, Kovalenko EV, et al. Biomarkers of myocardial and renal dysfunction in patients with coronary artery disease, diabetes mellitus and chronic kidney disease. *MIA Medical Bulletin*. 2023; 4(125): 52-56. Russian. DOI: 10.52341/20738080_2023_125_4_52
5. Zinellu A, Mangoni AA Cystatin C, COVID-19 severity and mortality: a systematic review and meta-analysis. *Journal of Nephrology*. 2022;35:59-68. DOI: 10.1007/s40620-021-01139-2
6. Orangereeva VN, Belaya OL, Kovalenko EV et al. The Capabilities of some modern biomarkers in predicting adverse cardiovascular events in patients with coronary heart disease and chronic kidney disease. *Effective pharmacotherapy*. 2024; 20 (31): 70-75. Russian. DOI 10.33978/2307-3586-2024-20-31-70-75

7. Liu F-T, Stowell SR The role of galectins in immunity and infection. *Nat Rev Immunol.* 2023; 23(8):479–494. DOI: 10.1038/s41577-022-00829-7
8. Rabkin SW, Tang JKK The utility of growth differentiation factor-15, galectin-3, and sST2 as biomarkers for the diagnosis of heart failure with preserved ejection fraction and compared to heart failure with reduced ejection fraction: a systematic review. *Heart Fail Rev.* 2021; 26(4):799–812. DOI: 10.1007/s10741-020-09913-3
9. Zhong X, Qian X, Chen G, et al. The role of galectin-3 in heart failure and cardiovascular disease. *Clin Exp Pharmacol Physiol.* 2019;46(3):197–203. DOI: 10.1111/1440-1681.13048
10. Rey JR, Caro-Codón J, Rosillo SO et al. Heart failure in COVID-19 patients: prevalence, incidence and prognostic implications. *Eur J of Heart Failure.* 2020; 22:2205–2215. DOI:10.1002/ehf.1990
11. Shi Q, Zhang X, Jiang F, et al. Clinical Characteristics and Risk Factors for Mortality of COVID-19 Patients With Diabetes in Wuhan, China: A Two-Center, Retrospective Study. *Diabetes Care.* 2020;43:1382–1391. DOI:10.2337/dc20-0598
12. Buliga-Finis ON, Ouatu A, Badescu MC, et al. Beyond the Cardiorenal Syndrome: Pathophysiological Approaches and Biomarkers for Renal and Cardiac Crosstalk. *Diagnostics.* 2022;12:773. DOI: 10.3390/diagnostics12040773
13. Galyavich AS, Tereshchenko SN, Uskach TM, et al. 2024 Clinical practice guidelines for Chronic heart failure. *Russian Journal of Cardiology.* 2024;29(11):6162. Russian. DOI: 10.15829/1560-4071-2024-6162
14. Ramos-Santos K, Cortes-Telles A, Uc-Miam ME et al. Cystatin C is a marker for acute kidney injury, but not for mortality among COVID-19 patients in Mexico. *Braz J Infect Dis.* 2022; 26(3):102365. DOI: 10.1016/j.bjid.2022.102365
15. Chen D, Sun W, Li J, et al. Serum Cystatin C and Coronavirus Disease 2019: A Potential Inflammatory Biomarker in Predicting Critical Illness and Mortality for Adult Patients. *Mediat Inflamm.* 2020; 2020: 3764515. DOI: 10.1155/2020/3764515
16. Zi M, Xu Y Involvement of cystatin C in immunity and apoptosis. *Immunol Lett.* 2018;196:80–90. DOI: 10.1016/j.imlet.2018.01.006
17. Anavi S, Tirosh O iNOS as a metabolic enzyme under stress conditions. *Free Radic Biol Med.* 2020; 146:16–35. DOI: 10.1016/j.freeradbiomed.2019.10.411
18. Karki R, Sharma BR, Tuladhar S, et al. Synergism of TNF- α and IFN- γ triggers inflammatory cell death, tissue damage, and mortality in SARS-CoV-2 infection and cytokine shock syndromes. *Cell.* 2021;184(1):149–168.e17. DOI: 10.1016/j.cell.2020.11.025
19. Liua Y, Xiaa P, Caob W et al. Divergence between serum creatinine and cystatin C in estimating glomerular filtration rate of critically ill COVID-19 patients. *RENAL FAILURE.* 2021; 43(1):1104–1114. DOI: 10.1080/0886022X.2021.1948428
20. Li Y, Yang S, Peng D, et al. Predictive value of serum cystatin C for risk of mortality in severe and critically ill patients with COVID-19. *World J Clin Cases.* 2020; 8(20): 4726–4734. DOI: 10.12998/wjcc.v8.i20.4726 90
21. Molvin J, Jujic A, Bachus E, et al. Cardiovascular biomarkers predict post-discharge re-hospitalization risk and mortality among Swedish heart failure patients. *ESC Heart Fail.* 2019; 6(5):992–999. DOI: 10.1002/ehf2.12486
22. Méndez AB, Azancot MA, Olivella A et al. New aspects in cardiorenal syndrome and HFpEF. *Clin Kidney J.* 2022;15(10):1807–1815. DOI: 10.1093/ckj/sfac133
23. Polat V, Bozcali E, Uygün T et al. Diagnostic significance of serum galectin-3 levels in heart failure with preserved ejection fraction. *Acta Cardiol.* 2016;71(2):191–197. DOI: 10.2143/AC.71.2.3141849
24. Kanukurti J, Mohammed N, Sreedevi NN, et al. Evaluation of Galectin-3 as a Novel Diagnostic Biomarker in Patients with Heart Failure with Preserved Ejection Fraction. *J Lab Physicians.* 2020;12(2):126–132. DOI: 10.1055/s-0040-1716608
25. Pecherina T, Kutikhin A, Kashtalap V, et al. Serum and Echocardiographic Markers May Synergistically Predict Adverse Cardiac Remodeling after ST-Segment Elevation Myocardial Infarction in Patients with Preserved Ejection Fraction. *Diagnostics (Basel).* 2020;10(5):301. DOI: 10.3390/diagnostics10050301
26. Roberts MA, Srivastava PM, Hare DL, et al. Effect of haemodialysis and residual renal function on serum levels of galectin-3, B-type natriuretic peptides and cardiac troponin T. *Nephrology (Carlton).* 2018; 23(12):1131–1138. DOI: 10.1111/nep.13179
27. Horiuchi YU, Wettersten N, VAN Veldhuisen DJ, et al. Galectin-3, Acute Kidney Injury and Myocardial Damage in Patients With Acute Heart Failure *J Card Fail.* 2023; 29(3):269–277. DOI: 10.1016/j.cardfail.2022.09.017
28. Edelmann F, Holzendorf V, Wachter R, et al. Galectin-3 in patients with heart failure with preserved ejection fraction: results from the Aldo-DHF trial. *Eur J Heart Fail.* 2015;17(2):214–223. DOI: 10.1002/ehf.203
29. Hasenfuss G, Pieske B, Wachter R, et al. The diagnostic and prognostic value of galectin-3 in patients at risk for heart failure with preserved ejection fraction: results from the DIAST-CHF study. *ESC Heart Fail.* 2021; 8(2): 829–841. DOI: 10.1002/ehf2.13174

Clinical significance of 24-hour blood pressure monitoring in patients with ankylosing spondylitis

Yupatova M.I., Nikitina N.M., Gafurova L.R.

Saratov State Medical University named after V.I. Razumovsky,
Ministry of Health of the Russian Federation, Saratov, Russia.

AUTHORS

Maria I. Yupatova, Assistant Professor, Department of Hospital Therapy, Faculty of Medicine, Saratov State Medical University named after V.I. Razumovsky, Ministry of Health of the Russian Federation, Saratov, Russia. ORCID: 0000-0001-8120-4424

Natalia M. Nikitina, MD, PhD, Associate Professor, Head of the Department of Hospital Therapy, Faculty of Medicine, Saratov State Medical University named after V.I. Razumovsky, Ministry of Health of the Russian Federation, Saratov, Russia. ORCID: 0000-0002-0313-1191

Leila R. Gafurova, Assistant Professor, Department of Hospital Therapy, Faculty of Medicine, Saratov State Medical University named after V.I. Razumovsky, Ministry of Health of the Russian Federation, Saratov, Russia. ORCID: 0000-0001-5247-5815

The aim of the study was to determine the clinical significance of 24-hour ambulatory blood pressure monitoring (ABPM) in patients with ankylosing spondylitis (AS).

Methods. A cross-sectional, single-center study included 47 men with a confirmed diagnosis of AS, aged 18 to 60 years (mean age 42.51 ± 6.04 years), with a mean disease duration of 17.26 ± 9.03 years. All patients were divided into two groups depending on the presence of hypertension based on office blood pressure (BP) measurement. AS activity was assessed using BASDAI and ASDAS-CRP indices. ABPM was performed in all patients.

Results. Low AS activity was observed in 2 (4.3%) patients, moderate in 20 (42.6%), and high in 25 (53.2%). A normal circadian BP profile ("dipper" type) was found in 16 (34%) patients, while impaired circadian BP regulation was observed in 31 (66%). No significant differences were identified between the two groups regarding the prevalence of abnormal BP profiles ($p = 0.547$). The duration of AS was significantly longer in patients with arterial hypertension (AH) (both previously diagnosed and newly detected by ABPM) compared with those without AH according to ABPM data ($p = 0.035$). Masked AH, as revealed by ABPM, was found in 10 (43.5%) patients from

the second group; among them, 5 (50%) also had an impaired circadian BP profile.

Conclusion. Arterial hypertension in patients with spondyloarthritis may remain subclinical. Nearly half of the patients who denied having AH were found to have masked hypertension, and half of them exhibited abnormal circadian BP profiles. These findings underscore the necessity of ABPM for the early detection of hypertension in patients with AS.

Keywords: spondyloarthritis, ankylosing spondylitis, arterial hypertension, masked arterial hypertension, circadian blood pressure profile, cardiovascular comorbidity.

Conflict of interests: none declared.

Received: 25.06.2025

Accepted: 07.08.2025



For citation: Yupatova M.I., Nikitina N.M., Gafurova L.R. Clinical significance of 24-hour blood pressure monitoring in patients with ankylosing spondylitis. 2025. 13(47):26-30. DOI: 10.24412/2311-1623-2025-47-33-38

Introduction

Cardiovascular diseases (CVD) have been the leading cause of mortality worldwide for many years [1].

The development of CVD is promoted by risk factors (RF), which are classified into non-modifiable (sex, age, race, genetics, etc.) and modifiable ones (smoking, physical inactivity, arterial hypertension (AH), dyslipidemia, obesity, diabetes mellitus, etc.) [2].

A pooled analysis of data from 112 cohort studies conducted in 34 countries including 1,518,028 participants demonstrated that modifiable RF account for about 50% of CVD burden in the general population. Among them, AH is the most significant [2].

Among patients with rheumatic diseases (RD), CVD also rank first in the structure of mortality, while the risk of their development substantially exceeds that in the general population [3]. This is explained by the higher prevalence of traditional RF as well as factors associated with chronic inflammation itself [4].

According to the largest cross-sectional observational study of comorbidities and RF in patients with spondyloarthritis (SpA) (COMOSPA), cardiovascular RF are highly prevalent in this group, with AH being the most common [5].

Ankylosing spondylitis (AS) is a chronic inflammatory disease from the SpA group, characterized by obligatory involvement of the sacroiliac joints and/or spine, potentially progressing to ankylosis, with frequent involvement of entheses and peripheral joints [6]. The disease mainly affects young, working-age individuals, gradually leading to disability. The development of AH and, consequently, the acceleration of CVD reduce working capacity and significantly impair quality of life in this patient population.

Diagnosis of AH includes office blood pressure (BP) measurement and ambulatory blood pressure monitoring (ABPM).

Depending on the circadian BP profile, patients with AH can be divided into four groups. A normal profile is the “dipper” type, characterized by a 10–20% decrease in nocturnal BP compared with daytime values. Other profiles — non-dipper, night-picker, and over-dipper — are associated with an increased risk of cardiovascular complications [7].

ABPM also allows for the detection of masked hypertension (MH), which is diagnosed in patients with normal BP values measured in a clinical setting but elevated BP outside the clinic. MH was first described more than 20 years ago and immediately became a subject of scientific interest [8]. Over the years, it has been shown that the risk of complications in MH is higher than in normotensive individuals and comparable to the risk of fatal cardiovascular events in manifest AH [9].

In the general population, the prevalence of MH is 8.5–16.6% [10]. In RA it reaches 38.7% [11]. However, no such studies have been conducted in patients with AS, and thus the value of ABPM in this population remains undefined.

The aim of the study was to determine the clinical significance of 24-hour ABPM in patients with ankylosing AS.

Methods

A cross-sectional single-center study included 47 men with a confirmed diagnosis of AS (according to the New York criteria, 1984 [12]) who were treated in the rheumatology department of the Saratov Regional Clinical Hospital. The mean age of patients

was 42.51 ± 6.04 years, and the mean duration of AS was 17.26 ± 9.03 years. Low AS activity was observed in 2 patients (4.3%), moderate in 20 patients (42.5%), and high in 25 patients (53.2%).

All patients were divided into two groups depending on the presence of AH: Group 1 included patients with established AH ($n = 24$, 51.1%), and Group 2 included patients in whom hypertension had not been previously diagnosed based on office BP measurement ($n = 23$, 48.9%). Patients in Group 1 were receiving adequate antihypertensive therapy at a stable dose for at least two weeks, maintaining BP within target values. The groups were comparable in age and AS activity ($p > 0.05$).

AS activity was assessed using the BASDAI (Bath Ankylosing Spondylitis Disease Activity Index) and ASDAS-CRP (Ankylosing Spondylitis Disease Activity Score) [12]. ABPM was performed using the BPLab device (LLC "Petr Telegin," Russia).

Exclusion criteria were the presence of other RD besides AS, thyroid dysfunction, previous myocardial infarction or stroke, as well as any comorbid pathology in the acute phase.

The main limitation of the study was the relatively small sample size, which did not allow us to establish associations between AS severity and the frequency of MH, as well as gender differences. The study had no sources of funding.

Statistical analysis

Statistical analysis was performed using StatTech v.4.8.11 software (developer – StatTech LLC, Russia).

Quantitative variables were assessed for normality using the Shapiro–Wilk test.

Quantitative variables with distributions consistent with normality were described using arithmetic means (M) and standard deviations (SD). The bound-

aries of the 95% confidence interval (95% CI) were provided as a measure of representativeness for mean values.

If the distribution of quantitative variables deviated from normality, the data were described using the median (Me) and lower and upper quartiles (Q1–Q3).

Categorical data were described using absolute values and percentages. The 95% CI for proportions was calculated using the Clopper–Pearson method.

Comparison of two groups for a quantitative variable with normally distributed data and equal variances was performed using Student's t-test; in the case of unequal variances, Welch's t-test was applied.

Comparison of two groups for a quantitative variable with a non-normal distribution was carried out using the Mann–Whitney U-test.

Comparison of proportions in 2x2 contingency tables was performed using Fisher's exact test (when the minimum expected frequency was less than 10). As a quantitative measure of effect when comparing relative indicators, odds ratios (OR) with 95% CI were calculated. In cases of zero cell frequencies in contingency tables, the odds ratio was calculated with the Haldane–Anscombe correction.

Comparison of proportions in multi-way contingency tables was carried out using Pearson's χ^2 test.

Differences were considered statistically significant at $p < 0.05$.

All participants were thoroughly informed and provided written informed consent. All procedures were approved by the Ethics Committee of Saratov State Medical University named after V.I. Razumovsky, Ministry of Health of Russia.

Results

The characteristics of the examined patients are presented in Table 1. The groups did not differ in terms

Table 1. Clinical characteristics of patients with AS depending on the presence of AH

Parameters	Group 1 (AS + AH), n=24	Group 2 (AS with no AH), n=23	p
Age, M (SD)	44.08 (6.68)	40.87 (4.92)	0.068
Disease duration, M (SD)	19.96 (8.93)	14.43 (8.43)	0.035*
CRP, Me [IQR]	7.35 [3.30; 24.50]	15.00 [3.90; 28.65]	0.595
Low activity, abs. (%)	0 [0.0%]	2 (8.7%)	0.310
Moderate activity, abs. (%)	10 (41.7%)	10 (43.5%)	
High activity, abs. (%)	14 (58.3%)	11 (47.8%)	
Continuous NSAID therapy	23 (95.8%)	22 (95.7%)	1.000
DMARD therapy	10 (43.5%)	10 (41.7%)	1.000
Biologic agents therapy	9 (37.5%)	4 (17.4%)	0.193
GC use	18 (75%)	13 (56.5%)	0.227

Note. * — the differences are statistically significant ($p < 0.05$)

Table 2. ABPM parameters in patients with AS depending on the initial presence of AH

Parameters	Group 1 (AS + AH), n=24	Group 2 (AS with no AH), n=23	p
Mean daytime SBP, M (SD), mmHg	133.96 [12.50]	125.74 [15.17]	0.048*
Mean daytime DBP, M (SD), mmHg	85.67 [8.11]	84.78 [11.18]	0.759
Mean nighttime SBP, M (SD), mmHg	125.33 [17.86]	113.30 [16.86]	0.022*
Mean nighttime DBP, M (SD), mmHg	78.21 [12.07]	75.52 [10.89]	0.428
Impaired circadian BP profile	17 (70.8%)	14 (60.9%)	0.547
Mean heart rate, M (SD), bpm	72.79 [11.60]	71.91 [10.06]	0.783

Note. * — the differences are statistically significant ($p < 0.05$)

of age or AS activity; however, patients in Group 1 had a longer disease duration ($p < 0.035$).

No significant differences were found between the groups in terms of C-reactive protein (CRP) levels or AS activity. The number of patients receiving nonsteroidal anti-inflammatory drugs (NSAIDs) on a regular basis, as well as those on disease-modifying antirheumatic drugs (DMARDs) or biologic agents, also did not differ significantly between the groups.

The results of ambulatory blood pressure monitoring (ABPM) in the examined patients are presented in Table 2.

According to ABPM, a normal circadian BP profile ("dipper" type) was identified in 16 patients (34%), while an abnormal circadian BP profile was observed in 31 patients (66%). Among these, the "non-dipper" profile was found in 22 patients (71%), the "over-dipper" profile in 3 patients (9.7%), and the "night-picker" profile in 6 patients (19.3%). There were no significant differences between the two groups in the frequency of abnormal circadian BP profiles ($p = 0.547$).

Mean systolic BP (SBP) was significantly higher in Group 1 during both daytime ($p = 0.048$) and nighttime ($p = 0.022$). No significant differences between the groups were observed in mean diastolic BP (DBP), either during the day ($p = 0.759$) or at night ($p = 0.428$).

MH according to ABPM data was detected in 10 patients (43.5%) in Group 2, of whom 5 patients (50%) had an abnormal circadian BP profile.

Discussion

The increased prevalence of CVDs in patients with RDs may be explained by a combination of traditional RFs with inflammation, oxidative stress, and the use of DMARDs, biologic agents, glucocorticoids (GCs), and NSAIDs [4, 13].

In the search for an association between CRP levels and the development of AH, a cohort of 20,525 female health professionals in the United States aged over 45 years with normotension at baseline was fol-

lowed for a mean of 7.8 years, with incident hypertension cases recorded. In adjusted models, an increase in baseline CRP by 1 SD was associated with an adjusted relative risk of developing AH of 1.14, suggesting that enhanced inflammation clearly preceded the clinical manifestation of elevated blood pressure [14]. The fact that AH appears to be associated with autoimmune inflammation underscores the need to investigate its coexistence with various RDs.

In the present study, it was found that patients with longer disease duration of AS more frequently had AH, while nearly every second patient with AS who denied having hypertension was found to have MH. This likely makes a significant contribution to the increased incidence of CVD in this patient population.

The prevalence of MH in AS is considerably higher than in the general population and higher than in patients with rheumatoid arthritis (Table 3).

Table 3. Frequency of MH in different population groups

Group	Frequency of MH
General population, n=7700	8.5–16.6 %
Patients with RA, n=31	38.7 %
Patients with AS, n=23	43.5 %

Thus, there is a strong tendency toward studying the BP profile in patients with RDs, and the results obtained highlight the problem of underdiagnosis of AH as one of the key contributors to the elevated CVD risk in this nosological group.

It is noteworthy that AH is a modifiable RF and therefore may serve as a successful target for interventions aimed at preventing CVD and mortality. Consequently, performing ABPM in patients with AS is essential for the timely detection and management of AH.

Conclusion

SpA are a group of diseases with a proven cardiovascular risk. AH in patients with AS may remain subclinical. Almost half of the patients who denied having AH

were found to have MH, and among them, half had an abnormal circadian blood pressure profile. This highlights the necessity of ABPM for the early detection of hypertension in patients with AS, since timely diagnosis will allow optimization of therapy and thereby delay the development of CVD in this population.

References

- Boytsov SA, Pogoseva NV Cardiovascular prevention 2022. Russian national guidelines. Russian Journal of Cardiology. 2023;28(5):5452. Russian. 2023;28(5):5452. DOI: 10.15829/1560-4071-2023-5452
- Global Cardiovascular Risk Consortium, Magnussen C, Ojeda FM, et al. Global Effect of Modifiable Risk Factors on Cardiovascular Disease and Mortality. N Engl J Med. 2023;389(14):1273-1285. DOI: 10.1056/NEJMoa2206916
- Conrad N, Verbeke G, Molenberghs G et al. Autoimmune diseases and cardiovascular risk: a population-based study on 19 autoimmune diseases and 12 cardiovascular diseases in 22 million individuals in the UK. Lancet. 2022; 3; 400(10354):733-743. DOI: 10.1016/S0140-6736(22)01349-6
- Toussiro E. The Risk of Cardiovascular Diseases in Axial Spondyloarthritis. Current Insights. Front Med (Lausanne). 2021; 8;8:782150. DOI: 10.3389/fmed.2021.782150
- Llop M, Gratacós J, Moreno M, et al. Sex differential impact of comorbidities in spondyloarthritis: data from COMOSPA study. RMD Open. 2024; 30;10(1):e003776. DOI: 10.1136/rmdopen-2023-003776
- Gaydukova IZ, Mazurov VI. Ankylosing spondylitis. In: Clinical rheumatology. Guide for doctors. 3rd ed., revised. Mazurov V.I., ed. M.:Ye-noto, 2021:127-141. Russian.
- Hermida RC, Mojón A, Fernández JR, et al. Elevated asleep blood pressure and non-dipper 24h patterning best predict risk for heart failure that can be averted by bedtime hypertension chronotherapy: A review of the published literature. Chronobiol Int. 2023;40(1):63-82. DOI: 10.1080/07420528.2021.1939367
- Kyriakoulis KG, Kollias A, Stergiou GS. Masked hypertension: how not to miss an even more silent killer. Hypertens Res. 2023;46(3):778-780. DOI: 10.1038/s41440-023-01182-5
- Kobalava ZhD, Konradi AO, Nedogoda SV, et al. Arterial hypertension in adults. Clinical guidelines 2020. Russian Journal of Cardiology. 2020;25(3):3786. Russian. DOI: 10.15829/1560-4071-2020-3-3786
- Parati G, Ochoa JE White Coat and Masked Hypertension. Hypertension. 2024. 24: 144-160. DOI: 10.1016/B978-0-323-88369-6.00012-8
- Osipova IV, Starodubova YN. Prediction of Cardiovascular Diseases in Women With Rheumatoid Arthritis. Kardiologiya. 2024; 31;64(1):67-79. Russian. DOI: 10.18087/cardio.2024.1.n2616
- Moiseev S, Novikov P, Gulyaev S, et al. Ankylosing spondylitis: diagnostic challenges and efficacy of upadacitinib. Clin Pharmacol Ther. 2021;30(4):62-70. Russian DOI: 10.32756/0869-5490-2021-4-62-70
- Liew JW, Ward MM, Reveille JD, et al. Nonsteroidal Antiinflammatory Drug Use and Association With Incident Hypertension in Ankylosing Spondylitis. Arthritis Care Res (Hoboken). 2020; 72(11):1645-1652. DOI: 10.1002/acr.24070
- Caiazzo E, Sharma M, Rezig AOM, et al. Circulating cytokines and risk of developing hypertension: A systematic review and meta-analysis. Pharmacol Res. 2024; 200:107050. DOI: 10.1016/j.phrs.2023.107050

Further studies are needed to clarify the role of MH in cardiovascular risk stratification not only in patients with AS but also in those with other RDs.

Conflict of interests: none declared.

Features of acute coronary syndrome associated with anxiety and depression disorders

Poponina T.M.¹, Repin A.N.², Poponina Y.S.^{1,2}, Gunderina K.I.²

¹ Siberian State Medical University of the Ministry of Healthcare of the Russian Federation, Tomsk, Russia.

² Cardiology Research Institute, Federal State Budgetary Scientific Institution «Tomsk National Research Medical Center of the Russian Academy of Sciences», Tomsk, Russia

AUTHORS

Tatiana M. Poponina, MD, PhD, Professor, Professor of the Department of Cardiology, Siberian State Medical University of the Ministry of Healthcare of the Russian Federation, Tomsk, Russia. ORCID: 0000-0002-6900-6190

Alexey N. Repin, MD, PhD, Professor, Head of the Department of Ambulatory Cardiology, Cardiology Research Institute, Federal State Budgetary Scientific Institution «Tomsk National Research Medical Center of the Russian Academy of Sciences», Tomsk, Russia. ORCID: 0000-0001-7123-0645

Yuliya S. Poponina, MD, PhD, Associate Professor of the Department of Cardiology, Siberian State Medical University of the Ministry of Healthcare of the Russian Federation; Cardiologist of the Department of Emergency Cardiology, Cardiology Research Institute, Federal State Budgetary Scientific Institution «Tomsk National Research Medical Center of the Russian Academy of Sciences», Tomsk, Russia. ORCID: 0000-0001-8167-1778

Kseniya I. Gunderina, MD, Cardiologist of the Department of Arterial Hypertension, Cardiology Research Institute, Federal State Budgetary Scientific Institution «Tomsk National Research Medical Center of the Russian Academy of Sciences», Tomsk, Russia. ORCID: 0000-0003-0804-4334

Cardiovascular Diseases (CVDs) are a leading cause of death, with the highest mortality rates observed during exacerbations of Coronary heart disease (CHD). A high incidence of affective disorders has been demonstrated in patients after acute coronary syndrome (ACS), both in the inpatient and outpatient settings.

The aim of this review is to present and analyze data from national and international studies on the features

of ACS, associated with anxiety and depression disorders, published between 2016 and 2024.

Methods. The review was prepared using materials from peer-reviewed articles published on PubMed, eLIBRARY.RU, cyberleninka.ru, and the other resources from 2016 to 2024 were used.

Results. Along with well-established cardiovascular risk factors (RFs), such as arterial hypertension,

smoking, and obesity, stress, anxiety, and depression are also recognized as risk factors. Due to hemodynamic disturbances in ACS, ischemic brain injuries may occur. These can lead to severe conditions, such as vascular delirium, as well as less severe disorders manifesting as cognitive impairment, sleep disorders, anxiety and/or depression.

Conclusion. To date, there are no clear, joint cardiology and psychiatry guidelines on diagnosis and management of psychiatric disorders in patients with ACS. Psychological and pharmacological interventions can have a positive effect and should be considered for ACS patients with depression, anxiety, and stress. Prior to discharge, all patients should be screened for mental disorders using validated questionnaires and, if necessary, a psychiatric consultation.

Keywords: Acute coronary syndrome, myocardial infarction, stress, anxiety, depression, risk factors, delirium, sudden cardiac death, antidepressants.

Conflict of interests: none declared.

Accepted: 25.06.2025

Received: 02.08.2025



For citation: Poponina T.M., Repin A.N., Poponina Y.S., Gunderina K.I. Features of Acute Coronary syndrome associated with anxiety and depression disorders. 2025. 13(47):31-36. DOI: 10.24412/2311-1623-2025-47-39-45

Introduction

Cardiovascular diseases (CVDs) remain a leading cause of death, with the highest mortality rates observed during exacerbations of coronary heart disease (CHD) [1]. A high incidence of affective disorders has been demonstrated in patients after acute coronary syndrome (ACS), both in the inpatient and outpatient settings [2, 3]. Experts of the European Society of Cardiology (ESC) and the American Heart Association have noted an increasing impact of psychosocial factors, such as stress, depression, and anxiety on the development and progression of CVDs [4-6]. Due to hemodynamic disturbances in ACS, ischemic brain injuries may occur. These can lead to severe conditions, such as vascular delirium, as well as less severe disorders manifesting as cognitive impairment, sleep disorders, anxiety and/or depression [7-9].

The aim of this review is to present and analyze data from national and international studies on the features of ACS, associated with anxiety and depression disorders, published between 2016 and 2024.

Methods

The review was prepared using materials from peer-reviewed articles published between 2016 and 2024 on databases such as PubMed, eLIBRARY.RU, CyberLeninka, and other relevant scientific platforms.

Results

Along with well-established cardiovascular risk factors (RFs), such as arterial hypertension, smoking,

and obesity, stress, anxiety, and depression are also recognized as risk factors [10]. However, to date, there are no clear, joint cardiology and psychiatry guidelines on diagnosis and management of psychiatric disorders in patients with ACS. Thus, an update to the ESC guidelines on ACS from 2023 [10] contains only one paragraph dedicated to mixed anxiety-depressive disorder (MADD) in ACS (13.2.4. Psychological considerations), reflecting the correlation between depression, anxiety, and stress and worse outcomes. Psychological and pharmacological interventions may have a positive effect and should be considered for ACS patients with depression, anxiety, and stress. Prior to discharge, all patients should be screened for mental disorders using validated questionnaires and, if necessary, a psychiatric consultation [5, 10].

The incidence of stress, anxiety and depression disorders in cardiovascular disease

The meta-analysis of 15 international clinical studies revealed that anxiety, depression, and psychosocial stress was associated with a 28% CVD risk increase [11]. MADD was most frequently observed in patients after acute myocardial infarction (AMI), with chronic heart failure (CHF), after cerebral strokes, in patients with atrial fibrillation (AF), as well as following advanced treatment procedures: coronary artery bypass graft (CABG) surgeries and implantations of cardioverter-defibrillators [12-14]. Chronic depression symptoms were identified in 65% of individuals with AMI, with every fifth of them diagnosed with an MADD [15]. According to research data [16], in pa-

tients with AMI, symptoms of anxiety, depression, and post-traumatic stress disorder (PTSD) occur in 9–60%, 11–20%, and 12–19% of cases, respectively. Stress, anxiety, and depression precede the development of AMI in 6–25% of cases [17]. Thus, ESC experts concluded the negative impact of anxiety, depression, and PTSD in patients with CVDs [18]. Derisi M.M. et al. mention in their work a 25.55% increase in the frequency of mental disorders among patients with coronary heart disease (CHD) from 2000 to 2021 compared to the period 1984–2000 [19]. Stress, anxiety, and depression are found in 58%, 33%, and 31% of patients with CVD, respectively [20, 21].

Stress, anxiety, depression, and the risk of adverse cardiovascular complications

A meta-analysis of 17 prospective studies, which analyzed the condition of 39,038 patients with ACS, found a 25% increased risk of adverse cardiovascular events (CV events) in patients with ACS associated with anxiety. However, after adjustment for depression, anxiety was not statistically significantly associated with mortality risk ($p=0.37$) [22]. Possible reasons for the link between depression and increased frequency of adverse CV events included lack of adherence to a healthy lifestyle (smoking, low physical activity, etc.), medication and rehabilitation non-adherence. ACS can cause post-traumatic stress, and PTSD triggered by ACS, in turn, can increase the risk of subsequent adverse CV events and mortality in patients. Clinically significant symptoms of ACS-induced PTSD are associated with a doubling of the risk of death and/or ACS recurrence (relative risk 2.00; 95% confidence interval (CI) 1.69–2.37) in patients with ACS who have clinically significant PTSD symptoms compared to patients without PTSD symptoms [22].

Mujkanovic J. et al. studied all cases of death among residents of Denmark aged 18 to 90 years in 2010 by analyzing death certificates and autopsy protocols: out of 4.3 million residents, the authors analyzed 45,703 death cases, of which 6,002 were caused by sudden cardiac death (SCD) [23]. The incidence of SCD was 1.79–6.45 times higher among patients with mental disorders than in the general population and was statistically significantly dependent on age ($p < 0.001$ in all age groups). In analyzing all death certifi-

icates and autopsy reports, the authors paid special attention to the medications that had been prescribed to the deceased¹.

Delirium and adverse outcomes in acute myocardial infarction

Delirium is an acute condition characterized by confusion, acute disturbances in attention, consciousness, or thinking, developing over several hours or days due to an underlying illness or medication side effects. Delirium cannot be explained by an alternative diagnosis or a comatose state [24].

Forms, key signs, and symptoms of delirium

There are three forms of delirium: hypoactive, hyperactive, and mixed. The key signs and symptoms of delirium include impaired attention (97%), long-term memory deterioration (89%), short-term memory deterioration (88%), disorientation (76%), reduced visuospatial abilities (87%), speech disturbances (57%), and disturbances of perception (50%) [25]. The Montreal Cognitive Assessment (MoCA) and the Confusion Assessment Method for the Intensive Care Unit (CAM-ICU) are used for the detection of delirium. The incidence of delirium in patients with AMI ranges from 1.4% to 11%. Delirium not only prolongs hospitalization but also increases in-hospital mortality by 1.4–1.9 times [26, 27].

Risk factors for delirium development

Predisposing factors for delirium include: advanced age, male sex, central nervous system disorders (dementia, stroke, Parkinson's disease), visual or hearing impairments, alcohol intoxication, immobilization, and alcoholism accompanied by dehydration [24, 25, 27].

Initiating factors for delirium development include: use and/or withdrawal of narcotic, toxic, and other psychotropic substances; severe acute illness; urinary tract infection; hypoxemia; hyponatremia; shock; anemia; pain syndrome; bladder catheterization; surgical interventions; admission to the intensive care unit (ICU); and a high number of medical procedures. Pharmacological risk factors for delirium development include: narcotic analgesics (primarily morphine); psychotropic drugs (benzodiazepines, tri-

¹ Mujkanovic J, Warming PE, Kessing LV et al. Use of antidepressant medication linked to substantial increase in risk of sudden cardiac death. <https://www.escardio.org/The-ESC/Press-Office/Press-releases/Use-of-antidepressant-medication-linked-to-substantial-increase-in-risk-of-sudden-cardiac-death>

cyclic antidepressants, haloperidol, phenothiazines, Belladonna-derived drugs, antiparkinsonian agents); drugs with anticholinergic activity; certain antibiotics; codeine; cimetidine; captopril; dipyridamole; hypothiazide; digoxin; furosemide; isosorbide dinitrate; nifedipine; prednisolone; theophylline; and other drugs such as antihistamines, laxatives, non-steroidal anti-inflammatory drugs, H₂-histamine receptor blockers, and anticoagulants.

The incidence of delirium can range from 5.7% in AMI patients admitted to the ICU to 40% in patients who have undergone open-heart surgery. Delirium, as a clinical manifestation of acute brain dysfunction syndrome occurring against the background of myocardial infarction (MI), is a serious complication of the underlying disease with an unfavorable prognosis.

Efficacy and safety of antipsychotic drugs in patients with delirium and myocardial Infarction

The efficacy and safety of antipsychotic drugs (neuroleptics) in patients with delirium and MI have not been sufficiently studied to date. An analysis of nine observational studies found that the odds of developing AMI were 1.88 times higher (odds ratio [OR] 1.88, 95% CI 1.39, 2.54) in users of antipsychotics compared to individuals who did not take antipsychotics. A subgroup analysis revealed an OR of 2.48 (95% CI 1.66, 3.69) among patients with schizophrenia and an OR of 2.64 (95% CI 2.48, 2.81) among short-term (<30 days) antipsychotics users. The authors of this meta-analysis confirm an increased risk of MI in users of antipsychotic drugs, demonstrating an elevated and more pronounced risk of AMI in short-term users [28]. In a number of studies, male sex, a diagnosis of schizophrenia, and antipsychotic exposure periods of < 60 days were associated with a higher risk of MI [29]. In a large nationwide cohort of patients admitted to the

hospital, Park Y. et al. found a slight increase in the risk of death within one week of initiating treatment with haloperidol compared to atypical antipsychotics [30]. The increased risk was noted within the first four days and was no longer apparent by the 5th day of observation. These findings are consistent with a large body of data from previous outpatient studies demonstrating a higher risk of mortality associated with the initiation of typical versus atypical antipsychotics [30]. The results of this study suggest that atypical antipsychotics may be a less harmful treatment option for elderly patients with AMI [30].

Conclusion

CHD and mental disorders are leading causes of morbidity and mortality not only in Russia but worldwide. Due to the high prevalence of depression and mortality from MI and CHD globally, the observed association between depression, MI, and SCD has significant public health implications. Psychological and pharmacological interventions may have a positive effect and should be considered for patients with ACS associated with depression, anxiety, and stress. Prior to discharge, all patients should be screened for mental disorders using validated questionnaires and, if necessary, a psychiatric consultation. The timely detection, treatment, and prevention of anxiety and depression can significantly reduce the risk of MI and SCD. However, to date, there are conflicting data regarding the efficacy and safety of antipsychotic drugs in patients with AMI and delirium. In the future, there is a critical need to develop national guidelines for the diagnosis and treatment of comorbid mental disorders in patients with CVD for cardiologists, internists, general practitioners, psychiatrists, and other specialists.

Conflict of interest: none declared.

References

1. Vaduganathan M, Mensah GA, Turco J V et al. The Global Burden of Cardiovascular Diseases and Risk. A Compass for Future Health. *J Am Coll Cardiol.* 2022;80(25):2361–2371. DOI: 10.1016/j.jacc.2022.11.005
2. Belialov FI, Sumin AN, Garganeeva NP et al. Mental disorders and cardiovascular diseases: guidelines of the baikal psychosomatic association and the siberian psychosomatic association. *Complex Issues of Cardiovascular Diseases.* 2024;13(3S):222–253. Russian DOI: 10.17802/2306-1278-2024-13-3S-222-253
3. Nonka TG, Lebedeva EV, Repin AN Clinical features of coronary artery disease and 5-year survival of patients after myocardial infarction against the background of depressive disorders. *Therapeutic archive.* 2024;96(1):17–21. Russian DOI:10.26442/00403660.2024.01.202560
4. Thompson DR Which Females Have the Highest Rates of Depression and Anxiety Following Acute Coronary Syndrome? *Cardiology.* 2023;148(1):20–22. DOI: 10.1159/000528624

5. Visseren FLJ, Mach F, Smulders YM et al. 2021 ESC Guidelines on cardiovascular disease prevention in clinical practice. *Eur Heart J.* 2021;42(34):3227-3337. DOI:10.1093/eurheartj/ehab484
6. Levine GN, Cohen BE, Commodore-Mensah Y et al. Psychological Health, Well-Being, and the Mind-Heart-Body Connection: A Scientific Statement From the American Heart Association. *Circulation.* 2021;143(10):e763-e783. DOI: 10.1161/CIR.0000000000000947
7. Panteleev OO, Ryabov VV. Delirium in a patient with myocardial infarction. *The Siberian Journal of Clinical and Experimental Medicine.* 2022;37(3):49-55. Russian. DOI: 10.29001/2073-8552-2022-37-3-49-55
8. Wu Q, Kling JM. Depression and the Risk of Myocardial Infarction and Coronary Death. A Meta-Analysis of Prospective Cohort Studies. *Medicine.* 2016;95(6):e2815-9. DOI: 10.1097/MD.0000000000002815
9. Poponina TM, Gunderina KI, Poponina YuS, Ryabov VV. Analysis of the S100B protein level in patients with acute coronary syndrome associated with anxiety and depressive symptoms. *The Siberian Journal of Clinical and Experimental Medicine.* 2023;38(2):82-88. Russian DOI: 10.29001/2073-8552-2023-38-2-82-88
10. Byrne RA, Rossello X, Coughlan JJ, et al. 2023 ESC Guidelines for the management of acute coronary syndromes. *European Heart Journal.* 2023; 44: 3720-3826. DOI: /10.1093/eurheartj/ehad191
11. Gaffey AE, Gathright EC, Fletcher LM, Goldstein CM. Screening for Psychological Distress and Risk of Cardiovascular Disease and Related Mortality: A Systematized Review, Metaanalysis, and Case for Prevention. *J Cardiopulm Rehabil Prev.* 2022;42(6):404-415 DOI: 10.1097/HCR.0000000000000751
12. Pushkarev GS, Kuznetsov VA, Fisher YA, et al. Impact of Depressive Symptoms on All-Cause Mortality in Patients with Congestive Heart Failure After Cardiac Resynchronization Therapy. *Cardiology.* 2019;59(1):5-11. Russian DOI: 10.18087/cardio.2019.1.10211
13. Kronish IM, Edmondson D, Moise N, et al. Posttraumatic stress disorder in patients who rule out versus rule in for acute coronary syndrome. *Gen Hosp Psychiatry.* 2018 Jul-Aug;53:101-107. DOI: 10.1016/j.genhosppsych.2018.02.007
14. Ben-Shoshan J, Segman-Rosensteig Y, Arbel Y, et al. Comparison of Triggering and Nontriggering Factors in ST-Segment Elevation Myocardial Infarction and Extent of Coronary Arterial Narrowing. *Am J Card.* 2016;117(8):1219-1223. DOI: 10.1016/j.amjcard.2016.01.038
15. Tofler GH, Kopel E, Klempfner R, et al. Triggers and Timing of Acute Coronary Syndromes. *Am J Card.* 2017;119(10):1560-1565. DOI: 10.1016/j.amjcard.2017.02.022
16. Chan B, Buckley T, Tofler G, et al. Emotional Stress and Physical Exertion as Triggers of Acute Myocardial Infarction. *American Journal of Cardiology.* 2023;203:285-287. DOI: 10.1016/j.amjcard.2023.07.085
17. Smyth A, O'Donnell M, Lamelas P, et al. Physical Activity and Anger or Emotional Upset as Triggers of Acute Myocardial Infarction. *Circulation.* 2016;134:1059-1067. DOI: 10.1161/CIRCULATIONAHA.116.023142
18. Piepoli MF, Hoes AW, Agewall S et al. 2016 European Guidelines on cardiovas-cular disease prevention in clinical practice. *European Heart J.* 2016;37:2315-2381. DOI: 10.1093/eurheartj/ehw106
19. Derisi MM, Nasiri MJ, Aryan A et al. Prevalence of psychiatric disorders in patients with ischemic heart disease: A systematic review and meta-analysis. *J Res Med Sci.* 2022;27:12. DOI: 10.4103/jrms.JRMS_864_20
20. Karami N, Kazemini M, Karami A, et al. Global prevalence of depression, anxiety, and stress in cardiac patients: A systematic review and meta-analysis. *Journal of Affective Disorders.* 2023;324:175-189. DOI: 10.1016/j.jad.2022.12.055
21. Lissåker CT, Norlund F, Wallert J, et al. Persistent emotional distress after a first-time myocardial infarction and its association to late cardiovascular and non-cardiovascular mortality. *Eur J Prev Cardiol.* 2019;26(14):1510-1518. DOI: 10.1177/2047487319841475
22. Li J, Ji F, Song J, et al. Anxiety and clinical outcomes of patients with acute coronary syndrome: a meta-analysis. *BMJ Open.* 2020;10(7):e034135. DOI:10.1136/bmjopen-2019-034135
23. Mujkanovic J, Warming PE, Kessing LV et al. Nationwide burden of sudden cardiac death among patients with a psychiatric disorder. *Heart.* 2024; 110(23):1365-1371. DOI: 10.1136/heart-jnl-2024-324092
24. Patil S, Gonuguntala K, Rojulpote C et al. Delirium is an important predictor of mortality in elderly patients with ST-elevation myocardial infarction: Insight from National Inpatient Sample database. *Coron. Artery Dis.* 2020;31(8):665-670. DOI: 10.1097/MCA.0000000000000978
25. Grover S, Ghosh A, Sarkar S et al. Delirium in Intensive Care Unit: Phenomenology, subtypes, and factor structure of symptoms. *Indian J. Psychol. Med.* 2018;40(2):169-177. DOI: 10.4103/IJPSYM.IJPSYM_274_17
26. Abdullah A, Eigbire G, Salama A, et al. Impact of delirium on patients hospitalized for myocardial infarction: A propensity score analysis of the National Inpatient Sample. *Clin Cardiol.* 2018;41(7):910-915. DOI: 10.1002/clc.22972
27. Jäckel M, Zotzmann V, Wengenmayer T, et al. Incidence and predictors of delirium on the intensive care unit after acute myocardial infarction, insight from a retrospective reg-

- 36 Poponina T.M., Repin A.N., Poponina Y.S., Gunderina K.I.
Features of acute coronary syndrome associated with anxiety and depression disorders
DOI: 10.24412/2311-1623-2025-47-39-45
-

- istry. Catheterization and Cardiovascular Interventions. 2021;6:1072-1081. DOI: 10.1002/ccd.29275
28. Yu ZH, Jiang HY, Shao L, et al. Use of antipsychotics and risk of myocardial infarction: a systematic review and meta-analysis. *Br J Clin Pharmacol*. 2016;82(3):624-32. DOI:10.1111/bcp.12985
29. Huang K-L, Fang C-J, Hsu C-C, et al. Myocardial infarction risk and antipsychotics use revisited: a meta-analysis of 10 observational studies. *Journal of Psychopharmacology*. 2017;31(12):1544-1555. DOI: 10.1177/0269881117714047
30. Park Y, Bateman BT, Kim DH, et al. Use of haloperidol versus atypical antipsychotics and risk of in-hospital death in patients with acute myocardial infarction: cohort study. *BMJ*. 2018;360. DOI: 10.1136/bmj.k1218

Modern trends in hypolipidemic therapy

Panakhova D.Z., Mirzabekova K.N.

Dagestan State Medical University, Makhachkala, Russian Federation.

AUTHORS

Diana Z. Panakhova, MD, PhD, assistant lecturer of the Department of Faculty Therapy, Dagestan State Medical University, Makhachkala, Russian Federation. ORCID: 0000-0003-0719-8980

Karina N. Mirzabekova, 6th-year student of the medical faculty, Dagestan State Medical University, Makhachkala, Russian Federation. ORCID: 0009-0001-8034-3374

Despite treatment with statins, patients with excessive levels of low-density lipoprotein cholesterol (LDL-C) and triglycerides have a relatively high risk of developing complications from cardiovascular diseases (CVDs).

For the past ten years, research of various hypolipidemic medications has been in constant development. Hypolipidemic medication is applied for treatment of patients with statin intolerance or inability to reach the recommended LDL-C level.

Inclisiran is a small interfering ribonucleic acid, targeted at the proprotein convertase subtilisin/kexin type 9 (PCSK9), exerts effects similar to monoclonal antibodies against PCSK9. Bempedoic acid, an ATP-citrate lyase enzyme inhibitor, is nowadays considered a promising option of statin-intolerant patients' treatment. Pemafibrate, an agonist of the peroxisome proliferator-activated receptor alpha, had demonstrated an exceptional risk/benefit ratio during the phase 2 of the clinical trials, nevertheless, the phase 3 was terminated due to ineffectiveness shown by the late interim analysis. High doses of icosapent ethyl, the ethylic ester of eicosapentaenoic acid, has a beneficial effect on the cardiovascular system.

For the time being, there still are not enough data on the use of novel medications for the primary prevention of CVDs. For faster integration into clinical practice, safety and effectiveness of these methods, as well as cost-efficiency, have to be demonstrated by the randomized study of cardiovascular outcomes.

Keywords: Monoclonal antibodies to proprotein convertase subtilisin/kexin 9, inclisiran, bempedoic acid, pemafibrate, eicosapentaenoic acid, docosahexaenoic acid.

Conflict of interests: None declared.

Received: 30.06.2025

Accepted: 15.08.2025



For citation: Panakhova D.Z., Mirzabekova K.N. Modern hypolipidemic pharmacotherapy trends. International Heart and Vascular Disease Journal. 2025. 13(47):37-47. DOI: 10.24412/2311-1623-2025-47-46-58

Introduction

Cardiovascular diseases (CVDs) are a leading cause of mortality not only in Russia, but worldwide as well [1]. In 2022 alone, CVDs caused 831 557 deaths in the Russian Federation. The mortality from neoplasia was almost 3 times lower. Most deaths were associated with complications of coronary heart disease (CHD) (54.2% of all deaths) and cerebrovascular disease (29.2% of all deaths). These diseases are commonly caused by atherosclerosis, which is strongly associated with dyslipidemia¹.

The goal of hypolipidemic therapy is to reduce the risk of developing atherosclerotic CVDs, most commonly caused by elevated levels of low-density lipoprotein cholesterol (LDL-C). Statins are the first-line drugs for treating hypercholesterolemia. For patients at high risk of CVD, with familial hypercholesterolemia (FH), or with type 2 diabetes mellitus (T2DM), it is recommended to achieve a target LDL-C level of < 1.8 mmol/L and reduce it by ≥50% from the baseline [2]. Nevertheless, the persisting risk of CVDs demonstrates an ongoing clinical need for the development and implementation of medications that effectively lower the blood levels of atherogenic particles [3].

Sufficient data on the efficacy of novel lipid-lowering drugs and their potential for clinical application have been accumulated in the literature. This review includes an analysis of new hypolipidemic drugs that affect blood levels of LDL and triglycerides (TG).

Methods

All material for this review was compiled from peer-reviewed articles published over the past 10 years on platforms such as PubMed, eLIBRARY.RU, cyberleninka.ru and other research databases. This time frame was selected as it encompasses the publication of key clinical studies on the subject. Search terms included "hypolipidemic therapy," "PCSK9 inhibitors," "inclisiran," and "bempedoic acid."

Studies or articles that did not meet quality standards were excluded.

Results

LDL-C-lowering drugs

A number of clinical and genetic studies have demonstrated that LDL particles cause CVDs, and that reducing CVD risk is dependent on lowering blood LDL-C levels [4].

Inhibition of Proprotein Convertase Subtilisin/Kexin Type 9 (PCSK9) with monoclonal antibodies

Proprotein convertase subtilisin/kexin type 9 (PCSK9) is a liver enzyme that binds to LDL-C receptors on hepatocyte membranes and reduces their expression. As a result of its action, the level of LDL-C in the plasma increases.

Evolocumab and alirocumab are monoclonal antibodies that inhibit this enzyme. These antibodies reduce the concentration of PCSK9 in the blood plasma, leading to increased expression of LDL-C receptors and, consequently, a decrease in blood LDL-C levels. PCSK9 inhibitors also reduce TG and lipoprotein(a) levels while simultaneously increasing apolipoprotein A-I (apoA-I) levels [5].

Evolocumab is indicated for patients with heterozygous and homozygous FH. It can be used as monotherapy or in combination for patients already on statin therapy who are not achieving their target LDL-C levels. This drug is also recommended for patients with statin intolerance [6, 7]. In patients with homozygous familial hypercholesterolemia, evolocumab reduced plasma LDL-C levels by 31% [8]. The cardiovascular outcomes study FOURIER (Further cardiovascular Outcomes Research with PCSK9 Inhibition in subjects with Elevated Risk) confirmed the feasibility of using evolocumab for hyperlipidemia. This study followed 27,564 patients with a plasma LDL-C level ≥ 1.8 mmol/L or non-high-density lipoprotein cholesterol (non-HDL-C) ≥ 2.6 mmol/L, who had atherosclerotic changes in various vascular beds and were concurrently receiving treatment with statins and ezetimibe. According to the results of this study, adding evolocumab to therapy reduced LDL-C levels by 59% from 2.4 mmol/L to 0.78 mmol/L, and the risk of CVD was reduced by 15% (hazard ratio (HR) = 0.85; 95% confidence interval (CI): 0.79-0.92, p <0.001) [9, 10].

¹ Federal State Statistics Service (Rosstat) «Demographic Yearbook of Russia 2023» <https://rosstat.gov.ru/folder/210/document/13207>

In the prospective, randomized, double-blind, placebo-controlled EVOPACS study, serum LDL-C levels were measured in 308 patients hospitalized with acute coronary syndrome (ACS) and elevated LDL-C levels, who had been on statin therapy for at least 4 weeks prior to hospitalization (LDL-C ≥ 1.8 mmol/L with high-intensity statin therapy; ≥ 2.3 mmol/L with low- or moderate-intensity statin therapy; or ≥ 3.2 mmol/L in patients taking statins inconsistently). Patients were randomized 1:1 to receive either evolocumab 420 mg subcutaneously or placebo in the hospital and for an additional 4 weeks after discharge (totaling 8 weeks), in addition to atorvastatin 40 mg therapy. The mean LDL-C level decreased from 3.61 to 0.79 mmol/L at week 8 in the evolocumab group and from 3.42 to 2.06 mmol/L in the placebo group; the difference in the mean percentage change from baseline was $\sim 40.7\%$ [95% CI: 45.2 to 36.2%; $p < 0.001$]. An LDL-C level of < 1.8 mmol/L was achieved at week 8 in 95.7% of patients in the evolocumab group compared to 37.6% in the placebo group, confirming the feasibility of using evolocumab [11].

The benefits of PCSK9 inhibitor drugs are also demonstrated by the results of the multicenter, double-blind, placebo-controlled, randomized clinical trial GLAGOV [10]. The study included 968 patients aged 18 years and older with specific clinical indications identified during coronary angiography. Specifically, they had at least one epicardial coronary stenosis obstructing 20% or more of the vessel lumen. Furthermore, these patients had a target vessel that could be visualized and whose obstruction did not exceed 50%. At the start of the study, patients were on stable-dose statin therapy for at least 4 weeks and had an LDL-C level of ≥ 2.1 mmol/L or between 1.6 and 2.1 mmol/L with one major or three minor risk factors for CVD. Major risk factors included: non-coronary atherosclerotic vascular disease, myocardial infarction (MI), or hospitalization for unstable angina within the previous 2 years, as well as type 2 diabetes. Minor risk factors included: smoking, hypertension, low high-density lipoprotein cholesterol (HDL-C) levels, family history of early coronary heart disease, high-sensitivity C-reactive protein (hs-CRP) level ≥ 0.2 mmol/L, or age ≥ 50 years for men and ≥ 55 years for women. The study patients were randomized to receive monthly subcutaneous injections of either evolocumab 420 mg (484 patients) or placebo (484 patients) for 76 weeks, in addition to statins. The

primary efficacy measure was the change in percent atheroma volume (PAV) from baseline to week 78, measured by serial intravascular ultrasound (IVUS). Secondary efficacy measures were the change in total atheroma volume (TAV) and the percentage of patients with plaque regression.

The safety and tolerability of evolocumab were also assessed. Among the 968 patients treated, lower LDL-C levels were achieved in the evolocumab group: 2.41 mmol/L in the placebo group versus 0.95 mmol/L in the evolocumab group, with a between-group difference of -1.46 mmol/L [95% CI: -1.55 to -1.38 mmol/L; $p < 0.001$]. The primary efficacy parameter, PAV, increased by 0.05% with placebo and decreased by 0.95% with evolocumab — a difference of -1.0% [95% CI: -1.8% to -0.64% ; $p < 0.001$]. The secondary efficacy parameter, TAV, decreased by 0.9 mm³ with placebo and by 5.8 mm³ with evolocumab, a difference of -4.9 mm³ [95% CI: -7.3 to -2.5 mm³; $p < 0.001$]. Evolocumab induced plaque regression in a greater percentage of patients than placebo: 64.3% vs. 47.3%, a difference of 17.0% [95% CI: 10.4 to 23.6%]; for PAV: 61.5% vs. 48.9%, a difference of 12.5% [95% CI: 5.9 to 19.2%; $p < 0.001$]; for TAV, $p < 0.001$. Thus, it can be concluded that the combination of evolocumab and statins promotes the regression of atherosclerotic plaques in patients with coronary heart disease, as confirmed by IVUS data [12].

The efficacy and safety of another drug in this class, alirocumab, in patients with high cardiovascular risk and uncontrolled hypercholesterolemia despite maximally tolerated statin doses, were established in the multicenter, double-blind, randomized, placebo-controlled study ODYSSEY COMBO II. This study included 720 patients with confirmed CVD and elevated LDL-C ≥ 1.8 mmol/L, or without confirmed CVD history but with high cardiovascular risk and LDL-C ≥ 2.6 mmol/L, despite maximum statin doses (rosuvastatin 20/40 mg, atorvastatin 40/80 mg, or simvastatin 80 mg) used for more than 1 year. Patients were randomized into 2 groups. The first group received alirocumab 75 mg subcutaneously every 2 weeks (plus oral placebo tablets). The second group received oral ezetimibe 10 mg daily (plus subcutaneous placebo injections) on top of statin therapy. At week 24, the mean reduction in LDL-C from baseline was $50.6 \pm 1.4\%$ for alirocumab versus $20.7 \pm 1.9\%$ for ezetimibe (difference $29.8 \pm 2.3\%$; $p < 0.0001$); 77.0% of patients receiving alirocumab and 45.6% of patients receiving

ing ezetimibe achieved LDL-C levels <1.8 mmol/L ($p < 0.0001$). The mean achieved LDL-C level at week 24 was 1.3 ± 0.04 mmol/L for alirocumab and 2.1 ± 0.05 mmol/L for ezetimibe and was maintained until week 52. Alirocumab was generally well-tolerated, with no signs of an excess of treatment-emergent adverse events observed [13].

The efficacy of long-term (78 weeks) treatment with alirocumab was demonstrated in two multicenter, randomized, double-blind, placebo-controlled studies, ODYSSEY FH I (486 patients) and FH II (249 patients), in patients with heterozygous familial hypercholesterolemia (HeFH) and inadequate LDL-C control despite maximally tolerated statin therapy (rosuvastatin 20–40 mg, atorvastatin 40–80 mg, or simvastatin 80 mg) with or without other lipid-lowering drugs for at least 4 weeks prior to the study (6 weeks for fenofibrate; other fibrates such as clofibrate, gemfibrozil, bezafibrate, ciprofibrate, except fenofibrate, were not permitted). Patients with HeFH without a history of cardiovascular events, as well as patients who had experienced MI or ischemic stroke, were included in the study if their LDL-C level did not reach the target value, according to current guidelines, for primary (≥ 2.6 mmol/L) or secondary (≥ 1.8 mmol/L) prevention respectively. Study patients were randomized in a 2:1 ratio to receive alirocumab 75 mg every 2 weeks or placebo. The alirocumab dose was blindly increased to 150 mg at week 12 if the patient's LDL-C level at week 8 was ≥ 1.8 mmol/L. Throughout the study, patients continued taking statins and other lipid-lowering medications. The mean LDL-C level decreased from a baseline of 3.7 mmol/L to 1.8 mmol/L after treatment (-57.9% compared to placebo) at week 24 in patients randomized to the alirocumab group in the ODYSSEY FH I study, and from 3.5 mmol/L to 1.8 mmol/L (-51.4% compared to placebo) in the ODYSSEY FH II study ($p < 0.0001$). This reduction was maintained until week 78. An LDL-C level of <1.8 mmol/L (regardless of cardiovascular risk) was achieved at week 24 by 59.8% and 68.2% of patients receiving alirocumab in the ODYSSEY FH I and FH II studies, respectively [14].

The multicenter, randomized, double-blind, placebo-controlled study ODYSSEY OUTCOMES demonstrated an improvement in cardiovascular outcomes after ACS in 18,924 patients receiving high-intensity statin therapy with the addition of 75 mg alirocumab. At baseline, the mean LDL-C level in the study

patients was 2.38 ± 0.80 mmol/L. In the alirocumab group, the mean LDL-C level at 4 months, 12 months, and 48 months after randomization was 1.0 mmol/L, 1.2 mmol/L, and 1.7 mmol/L respectively; in the placebo group, the mean LDL-C level at 4 months, 12 months, and 48 months after randomization was 2.4 mmol/L, 2.5 mmol/L, and 2.7 mmol/L respectively. In an analysis conducted in the alirocumab group (which excluded values measured after alirocumab discontinuation and after blinded substitution of alirocumab with placebo), the mean LDL-C level at 4 months, 12 months, and 48 months was 0.98 mmol/L, 1.1 mmol/L, and 1.4 mmol/L respectively; these levels were on average 62.7%, 61.0%, and 54.7% lower than the corresponding levels in the placebo group. The primary composite endpoint (PCE) included: death from coronary heart disease, nonfatal MI, fatal or nonfatal ischemic stroke, or unstable angina requiring hospitalization. The median follow-up duration was 2.8 years. A PCE event occurred in 903 patients (9.5%) in the alirocumab group and 1052 patients (11.1%) in the placebo group (HR = 0.85; 95% CI: 0.78–0.93; $p < 0.001$). A total of 334 patients (3.5%) died in the alirocumab group and 392 patients (4.1%) in the placebo group (HR = 0.85; 95% CI: 0.73–0.98). Thus, it can be concluded that alirocumab is capable of not only reducing LDL-C levels but also reducing the risk of cardiovascular complications and death in patients after ACS [15].

The effect of alirocumab on coronary artery atherosclerosis in patients who underwent percutaneous coronary intervention (PCI) for acute MI was studied in the recently conducted multicenter, double-blind, placebo-controlled, randomized clinical trial PACMAN-AMI. In this study, 300 patients who underwent PCI for acute MI (participant enrollment: from May 9, 2017, to October 7, 2020; final follow-up: October 13, 2021) were randomized to receive 150 mg of alirocumab (148 patients) subcutaneously every two weeks or placebo (152 patients). Treatment began less than 24 hours after urgent PCI and was administered for 52 weeks in addition to high-intensity statin therapy (rosuvastatin 20 mg). At 52 weeks, the mean change in PAV was -2.13% with alirocumab versus -0.92% with placebo, a difference of -1.21% (95% CI: -1.78% to -0.65% ; $p < 0.001$). The mean change in minimal fibrous cap thickness was 62.67 μm with alirocumab versus 33.19 μm with placebo, a difference of 29.65 μm (95% CI: 11.75–47.55; $p =$

0.001) [16]. Thus, in patients with acute MI, the addition of alirocumab to high-intensity statin therapy, compared with placebo, led to significantly greater regression of coronary plaque at 52 weeks.

Considering current clinical guidelines, PCSK9 inhibitors should be prescribed to patients receiving therapy with statins and ezetimibe at maximally tolerated doses, who have a high risk of CVD and a plasma LDL-C level ≥ 1.8 mmol/L [2].

With subcutaneous administration of alirocumab at doses of 75 and 150 mg and evolocumab at a dose of 140 mg every 2 weeks to patients with atherosclerotic CVD on background statin therapy, a 60% reduction in LDL-C level and a 15% reduction in the risk of CVD complications were observed [2].

PCSK9 inhibition via RNA interference

Inclisiran is a small interfering RNA that inhibits the synthesis of the PCSK9 protein in hepatocytes. The effect of this drug lasts longer than that of other lipid-lowering agents [17].

In the ORION-9 study evaluating the effect of inclisiran treatment on LDL-C levels in HeFH, which included 482 patients with HeFH and a baseline plasma LDL-C level ≥ 2.6 mmol/L on statin therapy, inclisiran sodium at a dose of 300 mg (equivalent to 284 mg of inclisiran free acid) was administered on days 1, 90, 270, and 450. The mean baseline LDL-C level was 4.0 ± 1.4 mmol/L. On day 510, inclisiran reduced plasma LDL-C levels by 47.9%. The mean absolute change in LDL-C level from baseline on day 510 was a reduction of 1.5 mmol/L (95% CI: -1.7 to -1.4 mmol/L) in the inclisiran group and an increase of 0.3 mmol/L (95% CI: 0.1 to 0.4 mmol/L) in the placebo group, with a between-group difference of 1.8 mmol/L (95% CI: -2.0 to -1.6 mmol/L; $p < 0.001$). A higher frequency of injection site reactions was noted in the inclisiran group compared to the placebo group (17.0% vs. 1.7%). Most adverse events were assessed as mild, with no serious complications identified [18].

In the ORION-10 study, which involved a total of 1561 patients with atherosclerotic CVD such as coronary heart disease, peripheral artery atherosclerosis, etc., a dose of 284 mg of inclisiran (as the free acid) was administered on days 1, 90, 270, and 450. The mean baseline LDL-C level was 2.71 ± 0.99 mmol/L. On day 510, inclisiran reduced plasma LDL-C levels by 52.3% (95% CI: 48.8 to 55.7%). The absolute

change in LDL-C level on day 510 was -0.05 mmol/L in the placebo group and -1.45 mmol/L in the inclisiran group, with a between-group difference of -1.40 mmol/L (95% CI: -1.48 to -1.32 mmol/L; $p < 0.001$). The percentage change on day 510 was 13.5% with placebo and -69.8% with inclisiran, representing a between-group difference of -83.3% (95% CI: -89.3% to -77.3% ; $p < 0.001$) [19].

Russian clinical guidelines approve inclisiran for subcutaneous administration at 284 mg, with a regimen of three doses in the first year and two doses annually thereafter [2].

Bempedoic Acid

ATP-citrate lyase (ACLY) catalyzes the ATP-dependent conversion of citrate and coenzyme A (CoA) into oxaloacetate and acetyl-CoA. Acetyl-CoA, a precursor of 3-hydroxy-3-methylglutaryl-CoA (HMG-CoA), is crucial for cholesterol biosynthesis. Thus, inhibition of ACLY leads to reduced levels of acetyl-CoA and cholesterol synthesis [20].

Bempedoic acid (BA) is a prodrug that acts as a selective ACLY antagonist after activation by acyl-CoA synthetase-1. Since this enzyme is primarily expressed in the liver, BA has minimal impact on other body tissues, particularly skeletal muscle. Unlike statins, patients treated with BA exhibited a lower incidence of myopathy, rhabdomyolysis, or other muscle-related adverse effects [20]. Among the most common side effects are increased plasma uric acid levels and the development of gout [21].

In addition to lowering LDL-C levels, BA also affects the level of high-sensitivity C-reactive protein (hs-CRP) [22].

The CLEAR Harmony and CLEAR Wisdom studies investigated the effect of BA therapy on cholesterol levels, involving 2230 and 779 patients, respectively, with atherosclerotic CVD and familial hypercholesterolemia and a plasma LDL-C level ≥ 1.8 mmol/L despite maximally tolerated statin therapy, with BA added to their regimen. Plasma LDL-C levels decreased by approximately 18%. In the CLEAR Harmony study, the baseline LDL-C was ≥ 1.8 mmol/L, while in CLEAR Wisdom, the mean baseline LDL-C was 3.1 ± 0.98 mmol/L. In CLEAR Harmony, at 12 weeks of BA therapy, the mean LDL-C level decreased by 0.5 mmol/L. Treatment with BA led to a significantly greater reduction in LDL-C levels compared to

the placebo group, both at week 12 ($p < 0.001$) and week 24 ($p < 0.001$). In the CLEAR Wisdom study, at week 12, BA reduced LDL-C significantly more than placebo (-15.1% vs. $+2.4\%$, respectively; $p < 0.001$). The mean LDL-C level at week 12 was 2.5 mmol/L in the BA group compared to 3.9 mmol/L in the placebo group [20, 23].

While the aforementioned studies assessed the efficacy of BA as an add-on lipid-lowering therapy in patients on maximum statin therapy, the CLEAR Serenity study included 345 patients with hypercholesterolemia and a history of intolerance to at least two statins (one statin at the lowest available dose). Patients were randomized 2:1 to receive oral BA 180 mg or placebo once a day for 24 weeks. At baseline, the mean LDL-C level was 4.08 ± 1.03 mmol/L, non-HDL-C was 4.99 ± 1.16 mmol/L, mean total cholesterol was 6.32 ± 1.2 mmol/L, apolipoprotein B (apoB) was 3.66 ± 0.81 mmol/L, and the median hs-CRP was 0.03 μ mol/L (Q1-Q3: 0.01-0.05 μ mol/L). BA treatment significantly reduced LDL-C from baseline to week 12 (placebo-corrected difference, -21.4% [95% CI: -25.1% to -17.7%]; $p < 0.001$). Significant reductions in BA group compared to placebo were also observed for non-HDL-C (-17.9%), total cholesterol (-14.8%), apoB (-15.0%), and hs-CRP (-24.3%); ($p < 0.001$ for all comparisons). BA was well-tolerated and safe. The most common muscle-related adverse event was myalgia, observed in 4.7% and 7.2% of patients receiving BA and placebo, respectively. This study expanded the evidence for the long-term use of BA in patients with statin intolerance [24].

BA affects blood levels of urea nitrogen, uric acid, and creatinine, increasing their levels. The drug also led to a decrease in hemoglobin levels. The incidence of gout was higher in the BA group compared to the placebo group (1.6 per 100 person-years vs. 0.5 per 100 person-years). The incidence of new-onset diabetes/hyperglycemia was lower in the BA group compared to the placebo group (4.7 per 100 vs. 6.4 per 100) [21].

The use of BA for the primary prevention of CVD was evaluated in the double-blind, randomized, placebo-controlled cardiovascular outcomes trial CLEAR Outcomes, involving 13,970 patients at high CVD risk, with statin intolerance, and an LDL-C level ≥ 2.59 mmol/L, who received BA (180 mg/day). Among the 13,970 patients included, 4,206 (30%) met the criteria for primary CVD prevention; of these, 2,100

were randomized to the BA treatment group and 2,106 to the placebo group. The mean LDL-C level was 3.69 mmol/L, mean HDL-C was 1.32 mmol/L, median TG was 4.19 mmol/L, and median hs-CRP was 0.06 mmol/L. Compared to placebo, BA reduced LDL-C by 21.3% after 6 months of treatment and reduced hs-CRP by 21.5% after 12 months of treatment. In the subgroup of high-risk CVD patients, the use of BA for primary CVD prevention was associated with a reduced incidence of major cardiovascular events. Specifically, BA reduced the risk of major adverse cardiovascular events by 13% (HR = 0.87; 95% CI 0.79–0.96; $p = 0.004$) [25, 26].

Thus, BA has become a long-awaited pharmacological alternative for lowering LDL-C levels both in patients not achieving target LDL-C levels on maximally tolerated statin therapy and in patients with statin intolerance. BA can be prescribed as monotherapy or as an add-on to treatment with maximally tolerated doses of statins and ezetimibe in patients who have not reached their target LDL-C level.

Agents for reducing triglyceride levels

Epidemiological and clinical studies clearly demonstrate a correlation between CVD risk and plasma TG levels. However, TGs play a secondary role in the formation of atherosclerotic plaques and are not directly involved in their development, as they can be broken down by most cells in the body. Consequently, TGs are not a direct cause of atherosclerosis. Nevertheless, recent attention has focused on the causal relationship between TGs, triglyceride-rich lipoproteins (TRLs), and their remnants in the risk of developing atherosclerosis. Genetic, epidemiological, and clinical studies suggest a causal link between TRLs and their remnants and an increased risk of atherosclerosis, and this dyslipidemia may be an independent risk factor for adverse cardiovascular events. Specifically, in patients with obesity, metabolic syndrome, diabetes, and chronic kidney disease, abnormal levels of TRLs and their remnants significantly increase the risk of developing atherosclerosis and CVD. The accumulation of excessively synthesized TRLs in the plasma, impaired function of enzymes involved in TRL lipolysis, and disrupted hepatic clearance of cholesterol-rich TRL remnants can lead to the deposition of TRLs and their remnants in the arteries, promoting the formation of foam cells and arterial wall inflammation. Thus, understanding the pathogenesis of

TRL-induced atherosclerosis and implementing targeted treatment may slow the progression of atherosclerosis, thereby reducing the morbidity and mortality from CVD, particularly coronary heart disease resulting from coronary artery atherosclerosis [27].

According to American clinical guidelines, omega-3 fatty acids or fibrates are prescribed to patients with hypertriglyceridemia for the prevention of pancreatitis [28]. European clinical guidelines recommend prescribing omega-3 polyunsaturated fatty acids in combination with statins for patients with elevated blood TG levels between 1.5 and 5.6 mmol/L [29].

Pemafibrate

Fibrates are TG-lowering drugs that act as agonists of the peroxisome proliferator-activated receptor alpha (PPAR α). These receptors are located on the membranes of hepatocytes, platelets, macrophages, as well as muscle, heart, kidney, and adipose tissue cells. Upon activation, PPAR α receptors stimulate or suppress genes responsible for metabolic processes in the body. A meta-analysis demonstrated that fibrates reduce the risk of CVD in a subgroup of patients with high baseline plasma TG levels [30].

Drugs in this class include: pemafibrate, bezafibrate, gemfibrozil, and fenofibrate.

Pemafibrate is the first selective PPAR α receptor agonist. Its activity exceeds that of fenofibrate, or more precisely its active form (fenofibric acid), by more than 2500 times. Pemafibrate demonstrates 2500 times greater affinity than fenofibric acid (the active metabolite of fenofibrate) for human PPAR- α , with over 5000 times greater affinity for PPAR- α than for PPAR- γ or - δ , thus requiring much lower doses and consequently limiting adverse effects [31].

The efficacy and safety of add-on therapy with pemafibrate for treating residual hypertriglyceridemia despite statin treatment were evaluated in a multicenter, randomized, double-blind, placebo-controlled, parallel-group, comparative clinical study on Japanese patients. The study included 751 patients with dyslipidemia who were taking pitavastatin. They were randomized into four groups of 188 patients each, assigned to receive either placebo or pemafibrate at doses of 0.1, 0.2, or 0.4 mg/day. These patients had residual dyslipidemia with mean fasting TG levels between 3.9 mmol/L and 4.3 mmol/L despite pitavastatin treatment, while the mean LDL-C level

ranged from 3.0 mmol/L to 3.2 mmol/L. Treatment continued for 12 weeks. Results from this first study showed that TG levels decreased significantly by approximately 50% from baseline to week 12 in all combination therapy groups, whereas no significant changes were observed in the pitavastatin monotherapy group. A dose-dependent reduction in TG levels was confirmed, and the decrease in TG levels from baseline to week 12 was statistically significant ($p < 0.001$) in all pemafibrate combination therapy groups compared to the pitavastatin monotherapy group [32].

The incidence of adverse events and adverse drug reactions in the pemafibrate combination therapy groups was comparable to that in the statin monotherapy groups, without any significant findings in the study [32].

The efficacy and safety of pemafibrate were also convincingly demonstrated among patients in a European randomized, double-blind, placebo-controlled Phase II study. The study included 408 patients who had been taking a stable dose of a statin (except pravastatin, lovastatin, or fluvastatin) for at least 12 weeks prior to the study and continued taking the same statin at an unchanged dosage throughout the study. At enrollment, the observed patients had fasting TG levels between 1.97 mmol/L and 5.65 mmol/L, HDL-C concentrations of ≤ 1.30 mmol/L for men and ≤ 1.43 mmol/L for women, and LDL-C concentrations not exceeding target values by more than 0.259 mmol/L. Patients were randomized to placebo or one of the following pemafibrate dosing regimens: 0.05 mg twice daily, 0.1 mg twice daily, 0.2 mg twice daily, 0.1 mg once daily, 0.2 mg once daily, or 0.4 mg once daily. Randomized patients received two tablets in the morning and two in the evening for 12 weeks. The tablets contained either the investigational drug (at doses of 0.05 mg, 0.1 mg, or 0.2 mg per tablet) or a matching placebo. Study results showed that pemafibrate reduced TG levels at any dosage ($p < 0.001$), with the greatest placebo-adjusted reduction from baseline to week 12 observed in the 0.2 mg twice-daily regimen group (54.4%). A significant increase in HDL-C levels, ranging from 7.4% to 12.9%, was also observed at all dosages except 0.1 mg once daily ($p < 0.05$). A significant increase in LDL-C levels, ranging from 9.2% to 20.5%, was observed at all dosages except 0.05 mg twice daily ($p < 0.05$). Pemafibrate was safe and generally well-tolerated. However, some pa-

tients exhibited a minor increase in serum creatinine and homocysteine concentrations [33].

Omega-3 Fatty Acid formulations

Omega-3 polyunsaturated fatty acids include eicosapentaenoic acid (EPA) and docosahexaenoic acid (DHA), which exert their effects indirectly through the activation of peroxisome proliferator-activated receptors (PPAR). By reducing the concentration of very-low-density lipoproteins, they lower triglyceride levels in the blood plasma [34].

Although omega-3 fatty acids are recommended for patients with severe hypertriglyceridemia in addition to statin therapy, the benefits of this approach for both primary and secondary prevention of CVD have not been confirmed by most conducted studies [34, 35].

For example, the multicenter, randomized, double-blind, placebo-controlled REDUCE-IT trial involved 8,179 patients with elevated plasma TG levels (1.69 to 5.63 mmol/L), LDL-C levels (1.06 to 2.59 mmol/L), and a history of atherosclerotic CVD or diabetes, who were receiving stable statin therapy for at least 4 weeks prior to the study. Patients were randomly assigned to groups receiving 2 grams of icosapent ethyl (an ethyl ester of eicosapentaenoic acid — an omega-3 polyunsaturated fatty acid ethyl ester) twice daily (total daily dose of 4 g) or placebo. Patients were followed for a median of 4.9 years. The primary composite endpoint was a combination of cardiovascular death, nonfatal MI, nonfatal stroke, coronary revascularization, or unstable angina. A key secondary endpoint (KSE) was the composite of cardiovascular death, nonfatal MI, or nonfatal stroke. PCE events occurred in 17.2% of patients in the icosapent ethyl (IE) group compared to 22.0% in the placebo group (HR=0.75; 95% CI: 0.68–0.83; $p<0.001$); corresponding rates for the KSE were 11.2% and 14.8% (HR=0.74; 95% CI: 0.65–0.83; $p<0.001$). The incidence of additional ischemic endpoints was significantly lower in the icosapent ethyl group than in the placebo group, including the rate of cardiovascular death — 4.3% vs. 5.2% (HR=0.80; 95% CI: 0.66–0.98; $p=0.03$). A greater number of patients in the icosapent ethyl group were hospitalized for atrial fibrillation or flutter compared to the placebo group (3.1% vs. 2.1%, $p=0.004$). Serious bleeding events were observed in 2.7% of patients in the

icosapent ethyl group and 2.1% in the placebo group ($p=0.06$) [36].

The effect of icosapent ethyl on the progression of coronary atherosclerosis in patients with elevated TG levels on statin therapy was studied in the randomized, double-blind, placebo-controlled EVAPORATE trial. This study included 80 patients aged 30 to 85 years with diagnosed coronary atherosclerosis ($\geq 20\%$ stenosis in one coronary artery by invasive coronary angiography or coronary computed tomography angiography (CCTA)), elevated fasting TG (1.52–5.63 mmol/L), and LDL-C levels from ≥ 1.04 to ≤ 2.98 mmol/L, who were on stable statin therapy (with or without ezetimibe), and adhering to a diet and exercise regimen for ≥ 4 weeks prior to the study. All patients followed a low-cholesterol diet and continued their current statin therapy during the study. Patients were randomized 1:1 to receive IE 4 g/day or mineral oil as a placebo to assess the rate of plaque volume progression via CCTA. Patients underwent interim scanning at 9 months and final scanning at 18 months using CCTA. The predefined PCE was the change in plaque volume at 18 months between the IE and placebo groups. The study results showed a significant reduction in the PCE, as IE reduced plaque volume by 17%, while in the placebo group, plaque volume more than doubled ($+109\%$) ($p=0.0061$). Significant differences in progression rates between the IE and placebo groups were also observed at the end of the study, including volumes of fibrous and fibrofatty plaques, which regressed in the IE group and progressed in the placebo group ($p<0.01$ for all). However, no statistically significant differences were found in the main lipid parameters (total cholesterol, LDL-C, HDL-C, and TG) from baseline to follow-up with either therapy [37].

However, opposing results were obtained in the randomized, double-blind, placebo-controlled, event-driven STRENGTH trial, which evaluated the long-term outcomes of risk reduction with omega-3 carboxylic acids in patients at high cardiovascular risk with hypertriglyceridemia on statin therapy [38, 39]. This drug is an omega-3 fatty acid formulation that undergoes an additional manufacturing step of hydrolyzing and distilling ethyl esters into omega-3 free fatty acids (omega-3 carboxylic acids) with a 75% concentration of EPA and DHA. Given its free fatty acid formulation, the drug does not require hydrolysis by pancreatic lipase, unlike conventional omega-3 ethyl

esters, to enhance intestinal absorption. This higher bioavailability is supported by its lack of dependence on dietary fat content.

The study included 13,086 patients aged ≥ 18 years who had been on a stable diet and taking statins for at least 4 weeks prior to the study initiation, with confirmed:

- LDL-C level < 2.59 mmol/L; however, subjects were also eligible for participation if their LDL-C level was ≥ 2.59 mmol/L, provided they were receiving high-intensity or maximally tolerated statin therapy, with or without ezetimibe;
- Atherogenic dyslipidemia, defined as TG levels ≥ 2.03 mmol/L and < 5.650 mmol/L and HDL-C < 1.09 mmol/L for men or < 1.22 mmol/L for women, measured after an overnight fast;
- High risk of future cardiovascular events.

The mean age of the studied patients was 62.5 ± 9.0 years. The cohort included 35% women and 65% men; 70% of all patients also had diabetes. The median LDL-C level was 1.94 mmol/L, the median TG level was 2.71 mmol/L, the median HDL-C level was 0.93 mmol/L, and the median hs-CRP level was 0.05 mmol/L.

Patients who met all inclusion criteria and voluntarily agreed to participate were randomized in a 1:1 ratio to receive either the study drug at 4 g/day or a corresponding placebo (corn oil). The study duration was 5 years.

The PCE was a composite of cardiovascular death, nonfatal MI, nonfatal stroke, coronary revascularization, and hospitalization for unstable angina. KSEs included:

- The composite of cardiovascular death, nonfatal MI, nonfatal stroke, coronary revascularization, and hospitalization for unstable angina in patients with established cardiovascular disease at baseline;
- The composite of cardiovascular death, nonfatal MI, and nonfatal stroke in the entire cohort and in patients with established cardiovascular disease at baseline;
- The composite of cardiac death, nonfatal MI, coronary revascularization, and hospitalization for unstable angina in the entire cohort and in patients with established cardiovascular disease at baseline;
- Cardiovascular death in the entire cohort and in patients with established cardiovascular disease at baseline;

- All-cause mortality in the entire cohort and in patients with established cardiovascular disease at baseline.

Prespecified tertiary efficacy endpoints (TCE) included newly diagnosed atrial fibrillation, thrombotic complications, and newly diagnosed heart failure. Changes in lipid levels, inflammatory markers, and EPA/DHA levels were also prespecified efficacy parameters. A retrospective analysis examined the association between plasma and erythrocyte EPA/DHA concentrations and the incidence of cardiovascular events [38, 39].

The study was terminated prematurely after 1,384 patients experienced a PCE, based on an interim analysis of the study results that indicated a low probability of clinical benefit for Epanova compared to the corn oil placebo. Of the 13,078 treated patients, 12,633 (96.6%) completed the study with confirmed PCE status. The PCE occurred in 785 patients (12.0%) receiving Epanova compared to 795 patients (12.2%) receiving placebo (HR = 0.99; 95% CI: 0.90–1.09; $p = 0.84$). The Epanova group had a higher incidence (24.7%) of gastrointestinal adverse effects (diarrhea, nausea, dyspepsia, and abdominal discomfort) compared to patients receiving placebo (14.7%) [38, 39].

Among patients at high cardiovascular risk taking statins, the addition of the omega-3 carboxylic acid formulation to usual background therapy did not lead to a significant change in the composite outcome of major adverse cardiovascular events compared to corn oil-based placebo. These results do not support the use of this specific omega-3 fatty acid formulation for reducing the incidence of major cardiovascular events in high-risk patients [38, 39].

Conclusion

Hypolipidemic therapy is primarily conducted to reduce the risk of CVDs. Statins have established themselves as first-line drugs for lowering plasma levels of LDL-C, a well-known risk factor for CVD. However, even with optimal statin therapy, a significant residual cardiovascular risk persists. Consequently, new drugs with mechanisms different from statins that lower LDL-C levels, as well as new targets beyond LDL-C for intervention to reduce the risk of atherosclerosis progression, are undoubtedly necessary.

Modern physicians have at their disposal a wide array of hypolipidemic drug classes with diverse mechanisms of action, the efficacy and safety of

which have been demonstrated in large, randomized, placebo-controlled trials. Consequently, for patients with dyslipidemia and atherosclerotic CVD who fail to achieve their target lipid profile on maximally tolerated statin therapy or in statin-intolerant patients, a

broad selection of alternative hypolipidemic agents is now available. It is crucial to emphasize, nonetheless, that statins remain the first-line therapy for initiating lipid-lowering treatment.

References

- Kosolapov VP, Yarmonova MV. The analysis of high cardiovascular morbidity and mortality in the adult population as a medical and social problem and the search for ways to solve it. *Ural medical journal*. 2021; 20 (1): 58-64. DOI: 10.52420/2071-5943-2021-20-1-58-64
- Ezhov MV, Arabidze GG, Drapkina OM, et al. Disorders of lipid metabolism. *Clinical Guidelines* 2023. *Russian Journal of Cardiology*. 2023;28(5):250-297. DOI: 10.15829/1560-4071-2023-5471
- François M, Konstantinos CK, et al. 2025 Focused Update of the 2019 ESC/EAS Guidelines for the management of dyslipidaemias: Developed by the task force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *European Heart Journal*. 2025; 00: 1-20. DOI: 10.1093/eurheartj/ehaf190
- Michaeli DT, Michaeli JC Established and Emerging Lipid-Lowering Drugs for Primary and Secondary Cardiovascular Prevention. *Am J Cardiovasc Drugs*. 2023;23(5):477-495. DOI: 10.1007/s40256-023-00594-5
- Moriarty PM, Thompson PD ODYSSEY ALTERNATIVE Investigators. Efficacy and safety of alirocumab vs ezetimibe in statin-intolerant patients, with a statin rechallenge arm: The ODYSSEY ALTERNATIVE randomized trial. *J Clin Lipidol*. 2015;9(6):758-769. DOI: 10.1016/j.jacl.2015.08.006.
- O'Donoghue ML, Giugliano RP, Wiviott SD, et al. Long-Term Evolocumab in Patients With Established Atherosclerotic Cardiovascular Disease. *Circulation*. 2022; 11;146(15):1109-1119. DOI: 10.1161/CIRCULATIONAHA.122.061620
- Stroes E, Colquhoun D, et al; GAUSS-2 Investigators. Anti-PCSK9 antibody effectively lowers cholesterol in patients with statin intolerance: the GAUSS-2 randomized, placebo-controlled phase 3 clinical trial of evolocumab. *J Am Coll Cardiol*. 2014; 17;63(23):2541-2548. DOI: 10.1016/j.jacc.2014.03.019
- Nissen SE, Stroes E; GAUSS-3 Investigators. Efficacy and Tolerability of Evolocumab vs Ezetimibe in Patients With Muscle-Related Statin Intolerance: The GAUSS-3 Randomized Clinical Trial. *JAMA*. 2016; 19;315(15):1580-90. DOI: 10.1001/jama.2016.3608
- Sabatine MS, Giugliano RP; FOURIER Steering Committee and Investigators. Evolocumab and Clinical Outcomes in Patients with Cardiovascular Disease. *N Engl J Med*. 2017; 4;376(18):1713-1722. DOI: 10.1056/NEJMoa1615664
- Raal FJ, Honarpour N, et al; TESLA Investigators. Inhibition of PCSK9 with evolocumab in homozygous familial hypercholesterolaemia (TESLA Part B): a randomised, double-blind, placebo-controlled trial. *Lancet*. 2015; 24;385(9965):341-50. DOI: 10.1016/S0140-6736(14)61374-X
- Koskinas KC, Windecker S, Pedrazzini G, et al. Evolocumab for Early Reduction of LDL Cholesterol Levels in Patients With Acute Coronary Syndromes (EVOPACS). *J Am Coll Cardiol*. 2019;19;74(20):2452-2462. DOI: 10.1016/j.jacc.2019.08.010
- Nicholls SJ, Puri R, Anderson T, et al. Effect of Evolocumab on Progression of Coronary Disease in Statin-Treated Patients: The GLAGOV Randomized Clinical Trial. *JAMA*. 2016; 13;316(22):2373-2384. DOI: 10.1001/jama.2016.16951
- Cannon CP, Cariou B; ODYSSEY COMBO II Investigators. Efficacy and safety of alirocumab in high cardiovascular risk patients with inadequately controlled hypercholesterolaemia on maximally tolerated doses of statins: the ODYSSEY COMBO II randomized controlled trial. *Eur Heart J*. 2015; 14;36(19):1186-94. DOI: 10.1093/eurheartj/ehv028
- Kastelein JJ, Ginsberg HN, Langslet G, et al. ODYSSEY FH I and FH II: 78 week results with alirocumab treatment in 735 patients with heterozygous familial hypercholesterolaemia. *Eur Heart J*. 2015; 14;36(43):2996-3003. DOI: 10.1093/eurheartj/ehv370
- Schwartz GG, Steg PG. ODYSSEY OUTCOMES Committees and Investigators. Alirocumab and Cardiovascular Outcomes after Acute Coronary Syndrome. *N Engl J Med*. 2018 Nov 29;379(22):2097-2107. DOI: 10.1056/NEJMoa1801174
- Räber L, Ueki Y; PACMAN-AMI collaborators. Effect of Alirocumab Added to High-Intensity Statin Therapy on Coronary Atherosclerosis in Patients With Acute Myocardial Infarction: The PACMAN-AMI Randomized Clinical Trial. *JAMA*. 2022; 10;327(18):1771-1781. DOI: 10.1001/jama.2022.5218
- Krzysztof D, Mariusz G, et al. Inclisiran – New hope in the management of lipid disorders? *Journal of Clinical Lipidology*. 2020; 14;1:16-27. DOI: 10.1016/j.jacl.2019.11.001
- Raal FJ, Kallend D, et al. Inclisiran for the Treatment of Heterozygous Familial Hypercholesterolemia. *N Engl J Med*. 2020; 382 (16): 1520-1530. DOI: 10.1056/NEJMoa1913805
- Ray KK, Wright RS, et al. Two Phase 3 Trials of Inclisiran in Patients with Elevated LDL Cholesterol. *N Engl J Med*. 2020; 382 (16): 1507-1519. DOI: 10.1056/NEJMoa1912387

20. Goldberg AC, Leiter LA, et al. Effect of bempedoic acid vs placebo added to maximally tolerated statins on low-density lipoprotein cholesterol in patients at high risk for cardiovascular disease: the CLEAR wisdom randomized clinical trial. *JAMA*. 2019;322(18):1780-1788. DOI: 10.1001/jama.2019.16585.
21. Bays HE, Banach M, Catapano AL, et al. Bempedoic acid safety analysis: Pooled data from four phase 3 clinical trials. *J Clin Lipidol*. 2020;14(5):649-659.e6. DOI: 10.1016/j.jacl.2020.08.009
22. Ray KK, Nicholls SJ; CLEAR OUTCOMES Committees and Investigators. Efficacy and safety of bempedoic acid among patients with and without diabetes: prespecified analysis of the CLEAR Outcomes randomised trial. *Lancet Diabetes Endocrinol*. 2024;12(1):19-28. DOI: 10.1016/S2213-8587(23)00316-9
23. Ballantyne CM, Banach M., et al. Long-term safety and efficacy of bempedoic acid in patients with atherosclerotic cardiovascular disease and/or heterozygous familial hypercholesterolemia (from the CLEAR Harmony Open-Label Extension Study). *Am J Cardiol*. 2022;174:1-11. DOI: 10.1016/j.amjcard.2022.03.020
24. Laufs U, Banach M, Mancini GBJ, et al. Efficacy and Safety of Bempedoic Acid in Patients With Hypercholesterolemia and Statin Intolerance. *J Am Heart Assoc*. 2019; 2;8(7):e011662. DOI: 10.1161/JAHA.118.011662
25. Albosta M, Grant JK, Michos ED. Bempedoic Acid: Lipid Lowering for Cardiovascular Disease Prevention. *Heart Int*. 2023;17(2):27-34. DOI: 10.17925/HI.2023.17.2.1
26. Nissen SE, Menon V, et al. Bempedoic Acid for Primary Prevention of Cardiovascular Events in Statin-Intolerant Patients. *JAMA*. 2023;330(2):131-140. DOI: 10.1001/jama.2023.9696.
27. Xu D, Xie L, et al. Triglyceride-rich lipoproteins and cardiovascular diseases. *Frontiers in endocrinology*. 2024;15:1409653. DOI:10.3389/fendo.2024.1409653
28. Patel SB, Wyne KL, Afreen S, et al. American Association of Clinical Endocrinology Clinical Practice Guideline on Pharmacologic Management of Adults With Dyslipidemia. *Endocrine Practice*. 2025; 31:236-262. DOI:10.1016/j.eprac.2024.09.016
29. Mach F, Baigent C, Catapano AL, et al. 2019 ESC/EAS Guidelines for the management of dyslipidaemias: lipid modification to reduce cardiovascular risk. *European Heart Journal*. 2020;41:111-188. DOI:10.1093/eurheartj/ehz455.
30. Yamashita S, Arai H, Yokote K, et al. Efficacy and Safety of Pemafibrate, a Novel Selective Peroxisome Proliferator-Activated Receptor α Modulator (SPPAR α): Pooled Analysis of Phase 2 and 3 Studies in Dyslipidemic Patients with or without Statin Combination. *Int J Mol Sci*. 2019; 6;20(22):5537. DOI: 10.3390/ijms20225537
31. Pradhan AD, Paynter NP, Everett BM, et al. Rationale and design of the Pemafibrate to Reduce Cardiovascular Outcomes by Reducing Triglycerides in Patients with Diabetes (PROMINENT) study. *Am Heart J*. 2018;206:80-93. DOI: 10.1016/j.ahj.2018.09.011
32. Arai H, Yamashita S, Yokote K, et al. Efficacy and safety of K-877, a novel selective peroxisome proliferator-activated receptor α modulator (SPPAR α), in combination with statin treatment: two randomised, double-blind, placebo-controlled clinical trials in patients with dyslipidaemia. *Atherosclerosis* 2017 Jun;261:144-152. DOI: 10.1016/j.atherosclerosis.2017.03.032
33. Ginsberg HN, Hounslow NJ, Senko Y, et al. Efficacy and Safety of K-877 (Pemafibrate), a Selective PPAR α Modulator, in European Patients on Statin Therapy. *Diabetes Care*. 2022 Apr 1;45(4):898-908. DOI: 10.2337/dc21-1288
34. ASCEND Study Collaborative Group, Bowman L, Mafham M, et al. Effects of n-3 fatty acid supplements in diabetes mellitus. *N Engl J Med* 2018; 18;379(16):1540-1550. DOI: 10.1056/NEJMoa1804989
35. Aung T, Halsey J, Kromhout D, et al. Omega-3 Treatment Trialists' Collaboration. Associations of Omega-3 Fatty Acid Supplement Use With Cardiovascular Disease Risks: Meta-analysis of 10 Trials Involving 77 917 Individuals. *JAMA Cardiol*. 2018; 1;3(3):225-234. DOI: 10.1001/jamacardio.2017.5205.
36. Bhatt DL, Steg PG. REDUCE-IT Investigators. Cardiovascular Risk Reduction with Icosapent Ethyl for Hypertriglyceridemia. *N Engl J Med*. 2019 Jan 3;380(1):11-22. DOI: 10.1056/NEJMoa1812792
37. Budoff MJ, Bhatt DL, Kinninger A, et al. Effect of icosapent ethyl on progression of coronary atherosclerosis in patients with elevated triglycerides on statin therapy: final results of the EVAPORATE trial. *Eur Heart J*. 2020 Oct 21;41(40):3925-3932. DOI: 10.1093/eurheartj/ehaa652
38. Nicholls SJ, Lincoff AM, Bash D, et al. Assessment of omega-3 carboxylic acids in statin-treated patients with high levels of triglycerides and low levels of high-density lipoprotein cholesterol: Rationale and design of the STRENGTH trial. *Clinical cardiology*. 2018;41(10):1281-1288. DOI:10.1002/clc.23055
39. Nicholls SJ, Lincoff AM, Garcia M, et al. Effect of High-Dose Omega-3 Fatty Acids vs Corn Oil on Major Adverse Cardiovascular Events in Patients at High Cardiovascular Risk: The STRENGTH Randomized Clinical Trial. *JAMA*. 2020;324(22):2268-2280. DOI:10.1001/jama.2020.22258

2025 ESC Congress HOT LINE sessions: key clinical studies

At the 2025 European Congress of Cardiology, the results of 42 international multicenter clinical trials were presented in the HOT LINE sessions. The studies addressed various areas of cardiology, including arterial hypertension, coronary heart disease, atrial fibrillation, lipid-lowering therapy, pharmacotherapy for heart failure, intensive care in cardiology, and healthcare management in cardiology. New texts for five clinical guidelines were also

presented. A summary report presents the results of the most interesting studies.

Keywords: clinical trials, meta-analyses, clinical guidelines, cardiovascular diseases.

Received: 02.08.2025

Accepted: 08.08.2025

From August 29 to September 1, 2025, the regular Congress of the European Society of Cardiology was held in Madrid (Spain), combined this year with the World Congress of Cardiology. The main theme of the event was “Global Health: Cardiology Without Borders”.

Congress participants reviewed 5 new texts of clinical guidelines:

- for the treatment of valvular heart diseases;
- for the treatment of myocarditis and pericarditis;
- for the treatment of cardiovascular diseases in pregnancy;
- targeted update of the 2019 guidelines for the treatment of dyslipidemia;
- consensus statement on mental health and cardiovascular diseases.

The full texts of these documents are available online at <https://www.escardio.org/Guidelines/Clinical-Practice-Guidelines>.

The results of 42 specially selected clinical studies were reported for the first time at 10 HOT LINE scientific sessions. The studies were devoted to various

areas of cardiology, including the management of patients with cardiac arrhythmias, chronic heart failure, hypertrophic cardiomyopathy, dyslipidemia, arterial hypertension, after myocardial infarction, coronary revascularization, venous thromboembolism, in the perioperative period.

The most interesting studies are summarized below.

The **POTCAST** study involved 1200 patients with a high risk of ventricular arrhythmias — with an implantable cardioverter defibrillator — and with an initial plasma potassium level of 4.3 mmol/l or lower. After randomization in a 1:1 ratio, treatment was administered to increase plasma potassium levels to 4.5–5.0 mmol/l using potassium supplements, a mineralocorticoid receptor antagonist, or both plus dietary advice, or standard care. At a median follow-up of 39.6 months, the composite of the primary endpoint events (documented sustained ventricular tachycardia or corresponding cardioverter-defibrillator shock, unplanned hospitalization for more than 24 hours for arrhythmia or heart failure, or death

from any cause) occurred in 22.7% of patients in the high-normal potassium group compared with 29.2% in the standard care group ($p=0.01$). The rate of hospitalization for hyperkalemia or hypokalemia was similar in both groups.

The therapeutic efficacy of the cardiac glycoside digitoxin in patients with heart failure and reduced ejection fraction was assessed in the **DIGIT-HF** study. A total of 1,212 patients with New York Heart Association functional class III or IV chronic heart failure with a left ventricular ejection fraction of 40% or less, or functional class II with a left ventricular ejection fraction of 30% or less were randomized in a 1:1 ratio to receive digitoxin (at an initial dose of 0.07 mg once daily) or placebo in addition to drug therapy prescribed in accordance with current clinical guidelines. At a median follow-up of 36 months, the primary endpoint (death from any cause or hospitalization for worsening heart failure) occurred in 39.5% of patients in the digitoxin group and 44.1% in the placebo group ($p=0.03$), with a trend toward a reduction in mortality (by 14%) in the cardiac glycoside group and a comparable incidence of serious adverse events (4.7% versus 2.8% of cases in the digitoxin and placebo groups, respectively).

Following the completion of the **VICTORIA** study ($n=5050$), the soluble guanylate cyclase stimulator vericiguat was approved for the treatment of advanced heart failure with reduced ejection fraction and received a Class IIb recommendation in European and North American guidelines. The subsequent VICTOR study ($n=6,105$) evaluated vericiguat in patients with heart failure with reduced ejection fraction without recent worsening. A pre-specified pooled analysis of patient-level data from the **VICTORIA** and **VICTOR** studies was conducted to determine the effect of vericiguat on clinical outcomes. Participants in both studies received current background heart failure therapy according to guidelines. The primary endpoint of cardiovascular death or hospitalization for heart failure occurred in 25.9% of 5,579 patients in the vericiguat group and 27.9% of 5,576 patients in the placebo group ($p=0.0088$), including significant reductions in the risk of cardiovascular death (by 11%; $p=0.020$) and hospitalization for heart failure (by 8%; $p=0.043$). Therefore, vericiguat may be used as an additional treatment option for selected patients with heart failure with reduced ejection fraction.

In the **ODYSSEY-HCM** study, the cardiac myosin inhibitor mavacamten, approved for the treatment of adult patients with symptomatic obstructive hypertrophic cardiomyopathy, was evaluated in the non-obstructive form. After randomization, 289 patients received mavacamten (starting dose of 5 mg/day adjusted to a maximum dose of 15 mg/day depending on left ventricular ejection fraction) and 291 received placebo. Over a 48-week follow-up period, there were no significant differences in the two primary endpoints: the mean change in peak oxygen consumption was 0.52 ml/kg/min in the mavacamten group and 0.05 ml/kg/min in the placebo group ($p=0.07$), and the change in quality of life according to the 23-item Kansas City Cardiomyopathy Questionnaire was 13.1 points versus 10.4 points ($p=0.06$). In patients with non-obstructive hypertrophic cardiomyopathy, mavacamten did not lead to a significantly greater improvement in peak oxygen consumption or reduction in symptoms compared with placebo.

The cardiac myosin inhibitor aficamten was compared with metoprolol in patients with obstructive hypertrophic cardiomyopathy in the MAPLE-HCM trial. After randomization, 88 patients received aficamten at a dose of 5 to 20 mg/day plus placebo, and 87 received metoprolol at a dose of 50 to 200 mg/day plus placebo. During 24 weeks of observation, the change in peak oxygen consumption (the primary endpoint) was 1.1 ml/kg/min in the aficamten group and -1.2 ml/kg/min in the metoprolol group ($p<0.001$). Patients receiving aficamten had significant improvements in New York Heart Association functional class, Kansas City Cardiomyopathy Questionnaire quality of life, left ventricular outflow tract pressure gradient, N-terminal pre-brain natriuretic peptide level, and left atrial volume index compared with patients receiving metoprolol. There were no significant differences in left ventricular mass index or adverse event rates between the two treatment groups.

The open-label, randomized **REBOOT-CNIC** trial compared the results of contemporary treatment (coronary reperfusion and secondary prevention) in patients with ST-segment elevation myocardial infarction and left ventricular ejection fraction greater than 40% with beta-blocker therapy ($n=4207$) versus no therapy ($n=4231$). With a median follow-up of 3.7 years, the primary endpoint (a composite of death from any cause, reinfarction, or hospitalization for heart failure) occurred at equal rates ($p=0.63$) in the

two groups; there was no difference in the rates of death from any cause, reinfarction, hospitalization for heart failure, or safety outcomes.

The open-label, randomized **BETAMI-DANBLOCK** study included patients with a history of myocardial infarction and a left ventricular ejection fraction of at least 40% who were randomized within 14 days of the event to receive long-term beta-blocker therapy (n=2,783) or no beta-blocker therapy (n=2,791). At a median follow-up of 3.5 years, the combined primary endpoint (death from any cause, myocardial infarction, unplanned coronary revascularization, ischemic stroke, heart failure, or malignant ventricular arrhythmia) occurred in 14.2% of patients in the beta-blocker group and 16.3% in the no-beta-blocker group (p=0.03), with no significant differences in the incidence of individual components or safety parameters between the groups. In this study, in patients with myocardial infarction and a left ventricular ejection fraction of at least 40%, beta-blocker therapy resulted in a reduced risk of death or major adverse cardiovascular events compared with no beta-blocker therapy.

The phase 3 BaxHTN study enrolled patients with sitting systolic blood pressure 140 mmHg to less than 170 mmHg despite treatment with two antihypertensive agents (uncontrolled hypertension) or three or more such agents (resistant hypertension), including a diuretic. After a 2-week placebo run-in period, patients with sitting systolic blood pressure 135 mmHg or greater were randomized to receive the selective aldosterone synthase inhibitor baxdrostat 1 mg (n=264), baxdrostat 2 mg (n=266), or placebo (n=264) once daily. After 12 weeks, the change from baseline in sitting systolic blood pressure (primary endpoint) was -14.5 mmHg with 1 mg baxdrostat, -15.7 mmHg with 2 mg baxdrostat, and -5.8 mmHg with placebo. The placebo-corrected difference was -8.7 mmHg with 1 mg baxdrostat and -9.8 mmHg with 2 mg baxdrostat (p<0.001 for both comparisons). Potassium levels greater than 6.0 mmol/L were recorded in 2.3% of patients in the 1 mg baxdrostat group, 3.0% in the 2 mg baxdrostat group, and 0.4% in the placebo group. In patients with uncontrolled or resistant hypertension, the addition of baxdrostat to background therapy resulted in a significant reduction in sitting systolic blood pressure after 12 weeks compared with placebo.

Olezarsen is an antisense oligonucleotide that inhibits the synthesis of apolipoprotein C-III, result-

ing in lower plasma triglyceride levels. The drug was evaluated in the randomized phase 3 Essence-TIMI 73b study in patients with moderate hypertriglyceridemia (triglycerides 150 to 499 mg/dL) and increased cardiovascular risk or severe hypertriglyceridemia (triglycerides ≥500 mg/dL) who were randomized 1:3 to receive monthly subcutaneous administration of olezarsen 50 mg (n=254), olezarsen 80 mg (n=766), or placebo (n=329). After 6 months of treatment, the mean change in triglyceride levels adjusted for placebo was -58.4% in the olezarsen 50 mg group (p<0.001) and -60.6% in the olezarsen 80 mg group (p<0.001). The incidence of serious adverse events was similar in all study groups.

The optimal duration of anticoagulation after venous thromboembolism in patients with a transient precipitating event (e.g., surgery, trauma, or immobilization) and concomitant persistent risk factors has not been determined. In the randomized HI-PRO study, patients with VTE following a transient trigger, at least 1 persistent risk factor, and receiving anticoagulation for at least 3 months were randomized to receive apixaban 2.5 mg twice daily (n=300) or placebo (n=300) for 12 months. During the follow-up period, the primary efficacy endpoint (first symptomatic recurrence of VTE) occurred in 1.3% of patients in the apixaban group and 10.0% in the placebo group (p<0.001). The primary safety endpoint (first major bleeding event according to the International Society on Thrombosis and Haemostasis criteria) occurred with comparable frequency: major bleeding occurred in 1 patient in the apixaban group and in no patients in the placebo group, minor clinically significant bleeding was observed in 4.8% of patients in the apixaban group and in 1.7% in the placebo group (p=0.06). Rare fatal outcomes were not associated with cardiovascular or hemorrhagic causes.

The **AQUATIC** study included patients with chronic coronary syndrome who had a stent implanted more than 6 months before inclusion in the study, had a high atherothrombotic risk and received continuous oral anticoagulant therapy. After randomization, once-daily aspirin 100 mg (n=433) or placebo (n=439) was administered in addition to anticoagulants. The study was terminated early on the recommendation of an independent data and safety monitoring board after a median follow-up of 2.2 years due to excess death from any cause in the aspirin group. The primary efficacy endpoint (a composite of cardiovas-

cular death, myocardial infarction, stroke, systemic embolism, coronary revascularization, or acute limb ischemia) occurred in 16.9% of patients in the aspirin group and in 12.1% in the placebo group ($p=0.02$), and death from any cause occurred in 13.4% versus 8.4% of cases, respectively ($p=0.01$). The key safety indicator (major bleeding) was recorded in 10.2% of patients in the aspirin group and in 3.4% in the placebo group ($p<0.001$).

The **NEO-MINDSET** study enrolled patients after successful percutaneous coronary intervention for acute coronary syndromes who were randomized within the first 4 days of hospitalization to stop aspirin and then receive monotherapy with a potent P2Y₁₂ inhibitor (ticagrelor or prasugrel) ($n=1712$) or dual antiplatelet therapy (aspirin and a potent P2Y₁₂ inhibitor) ($n=1698$) for 12 months. During this follow-up period, the primary end point events (death from any cause, myocardial infarction, stroke, or urgent revascularization) occurred in 7.0% of patients in the monotherapy group and 5.5% in the dual antiplatelet therapy group ($p=0.11$ for noninferiority). Major or clinically relevant non-major bleeding occurred in 2.0% of patients in the monotherapy group and 4.9% of patients in the dual antiplatelet therapy group, and stent thrombosis occurred in 12 and 4 patients in the matched groups, respectively. Among patients who underwent successful percutaneous coronary intervention for acute coronary syndrome, monotherapy with a potent P2Y₁₂ receptor inhibitor was non-inferior to dual antiplatelet therapy in preventing the combined risk of death and ischemic events at 12 months.

The **TARGET-FIRST** trial included patients who had successfully undergone complete coronary revascularization with implantation of advanced drug-eluting stents within 7 days of myocardial infarction and who completed 1 month of dual antiplatelet therapy without ischemic complications or major bleeding. After randomization, patients were assigned to receive either P2Y₁₂ receptor inhibitor monotherapy ($n=961$) or dual antiplatelet therapy ($n=981$) for 11 months. During follow-up, the primary composite

endpoint (death from any cause, myocardial infarction, stent thrombosis, stroke, or major bleeding type 3 or 5 according to the Bleeding Academic Research Consortium definition) occurred in 2.1% of patients in the P2Y₁₂ receptor inhibitor monotherapy group and 2.2% of patients in the dual antiplatelet therapy group ($p=0.02$ for noninferiority). The primary secondary endpoint (Bleeding Academic Research Consortium type 2, 3, or 5 bleeding) occurred in 2.6% of patients in the P2Y₁₂ receptor inhibitor monotherapy group and 5.6% in the dual antiplatelet therapy group ($p=0.002$ for superiority). Stent thrombosis was rare and its incidence was similar in both groups.

In this open-label, registry-based, cardiothoracic surgery center-based clinical trial, TACSI, patients undergoing coronary artery bypass grafting for acute coronary syndrome were randomized to receive ticagrelor and aspirin ($n=1104$) or aspirin alone ($n=1097$) for 1 year. During this follow-up period, the combined primary endpoint event (death, myocardial infarction, stroke, or repeat revascularization) occurred in 4.8% of patients in the ticagrelor plus aspirin group and 4.6% in the aspirin alone group ($p=0.77$). The key secondary endpoint was all adverse clinical outcomes, defined as the primary endpoint events plus major bleeding, and its incidence was 9.1% of cases in the ticagrelor plus aspirin group versus 6.4% in the aspirin alone group. Major bleeding occurred in 4.9% of patients in the ticagrelor plus aspirin group and 2.0% in the aspirin alone group. In patients undergoing coronary artery bypass grafting for acute coronary syndrome, adding ticagrelor to aspirin does not reduce the risk of death, myocardial infarction, stroke, or repeat coronary revascularization compared with aspirin alone for 1 year.

The next congress of the European Society of Cardiology is scheduled to be held in Munich, Germany, from 28 to 31 August 2026.

The material was prepared by **Professor Kanorsky S.G.** (Krasnodar, Russia), edited by **Professor Mamedov M.N.** (Moscow, Russia).

Author Guidelines

Manuscript publication rules
in the International heart and vascular disease journal

Edit from December, 2021

Disclaimer: The rules came into effect from December 2021. The rules describe the conditions of publication of manuscripts (articles) through the site <http://www.heart-vdj.com>. The editorial Board is ready to answer questions and help authors by e-mail: submissions.ihvdj@gmail.com.

The *International heart and vascular disease journal* has been published since 2013. It is official journal of the Cardioprogress Foundation. The target audience of this peer-reviewed journal is cardiologists and internal disease specialists. The journal is primarily focused on questions of epidemiology, prevention, and cardiac pharmacotherapy. It also publishes lectures and literature reviews on various problems of modern cardiology, reports on new diagnostic methods, and other information which is important for the practitioners.

The General criteria for the publication of articles in the International heart and vascular disease journal are the relevance, novelty of the material and its value in theoretical and/or applied aspects.

The languages of publications are Russian and English. Journal is peer-reviewed, with multistage editing. Editorial board is presented by the leading cardiologists from different countries and Russia.

International heart and vascular disease journal aims to ensure that its publications fulfill the requirements of international publishing standards, such as the Uniform Requirements for Manuscripts Submitted to Biomedical Journals: Writing and Editing for Biomedical Publication, by the International Committee of Medical Journal Editors, ICMJE (<http://www.icmje.org>), and the recommendations by the

Committee on Publication Ethics, COPE (<http://www.publicationethics.org.uk>).

All clinical trials should be performed and described in full accordance with the CONSORT standards (<http://www.consort-statement.org>), observational research — STROBE (<http://www.strobe-statement.org>), systematic reviews and meta-analyses — PRISMA (<http://www.prisma-statement.org>), diagnostic accuracy — STAR (<http://www.stard-statement.org>).

I. The International heart and vascular disease journal accepts the following manuscripts:

1) *Original papers* present the results of clinical studies. The word limit is 3.000 (including references, tables, and figure legends). The maximal number of references is 15. The structured abstract should contain 5 sections (**Aim, Material and Methods, Results, Conclusion, and Key words**), and be no longer than 300 words.

2) *Lectures*, or clinically oriented reviews, are written by experts in broader areas of medicine. Lectures could be focused on epidemiology, pathophysiology, diagnostics, treatment, and prevention. The word limit is 5.000 (including references, tables, and figure legends). The maximal reference number is 80. The unstructured abstract is no longer than 150 words.

3) *Literature reviews* are focused on more specific topics, compared to lectures. The word limit is 4.500 (including references, tables, and figure legends). The maximal reference number is 50. The unstructured abstract is up to 150 words.

4) *Clinical case* is a brief report on a complex diagnostic problem and its solution, or a description of

a rare clinical observation. The word limit is 600 (including references, tables, and figure legends). The maximal number of references is 5. No abstract is required.

5) *Clinical opinion* informs the readers on the topics of cardiovascular medicine and related disciplines. The word limit is 2.500 (including references, tables, and figure legends). The maximal number of references is 15.

The journal accepts for publication original phase 2, 3 and 4 clinical studies. Literature reviews should be based on sources not older than 5 years.

II. Information about the article, which includes the following sections, is combined into a single file "letter (cover)":

1) the manuscript is not under consideration in another edition has not been previously published contains a full disclosure of the conflict of interest all authors meet the criteria of authorship, it was read and approved the author (s) are responsible for the power of attorney submitted in the manuscript materials. 6) all contact information of the author responsible for correspondence information about previous publications of the authors on the same topic or pre-publication.

If the manuscript is a part of the thesis, it is necessary **to specify** the estimated terms of thesis defense.

The "letter of direction (accompanying)" should be made out on one or two sheets. Using the form of the official institution-at the choice of the author's team. In the address: "to The chief editor of the Russian cardiology journal, academician of RAS, Professor Oganov R. G.". The signatures of **all authors** should be placed at the bottom.

"Directional (cover) letter" is scanned. File format. jpeg attached as an additional file of the manuscript.

The absence of a letter or incomplete text of the letter (not containing the above items) is the basis for refusal to accept the manuscript for consideration.

III. Registration on the Website and information about the authors.

Any of the authors can submit an article to the journal. Usually it is the one who then conducts correspondence with the editorial office and to whose mail notification letters come (when submitting a manuscript through the site, you can choose to send notifications to all authors).

The author registers on the site, entering his full name. In the form to be filled in when submitting an article, all authors and all additional information (places of work, positions, academic titles, institutions, ORCID — all authors) are indicated.

If the author has several places of work, it is written: 1. "The name of the institution..." 2. "Name of institution..." The name of the institution is written in abbreviated form, for example, Moscow state University, Moscow. Brackets are not put.

How to fill in the article metadata: all data that is entered in the "article metadata" must exactly match the data specified in the text of the article!

Authors' names (you can not write in full, the format of the journal provides for the publication of names and initials. Therefore, in the "Windows", where the name and patronymic of the authors are written in capital letters with a dot (example: A.).

Names of institutions (write the official name. At the same time — there is a reduction of Federal, STATE, etc.; the quotation marks are placed; Ministry of health of Russia, a city without the letter G.

Positions and titles (using traditional abbreviations: PhD, senior researcher, leading researcher, PhD, C.b.N., MD), head reduces to the head., then write the full name of the laboratory/Department / Department; Director, head, Professor — is not reduced.

The order of the authors. Authors' priority should be entered into the system in accordance with the order of the article. The movements are made by small arrows "top" / "bottom", which are located under the data of each of the authors. The data of the author responsible for the correspondence, put a dot in a circle denoting this information. Other authors point do not put.

Summary. Sections of the abstract should exactly match the sections prescribed In the rules for authors. If the sections are not correct, the Editors will ask to correct them. What the authors are currently publishing on the site will then be included in all systems after the final publication. Be careful!

Making literary references. Submitted article will not be reviewed until the correction of literary references in accordance with the rules for authors is made. The authors "forget" and somewhere to remove point (such inconsistencies can be corrected in the Revision), but if the design literature is radically different from what is required or present hyperlinks,

the Editors will not start with the article to eliminate errors.

Keyword. They are written with a small letter, separated by a semicolon. At the end put a point. In the text of the article the keywords are written separated by commas.

A file is prepared separately in Word, which is then sent as an additional file. The file must contain:

Title page of the manuscript. The title of the manuscript is written in capital letters, without hyphenation, in bold. Initials and surnames of authors-Ivanov I. I., Petrov P. p. the full name of organization (s) from which (s) there was a manuscript, the city, the country is Given. Footnotes are in Arabic numerals after the authors' names and before the names of institutions.

Example of design:

THE PREVALENCE OF RISK FACTORS OF NONCOMMUNICABLE DISEASES IN THE RUSSIAN POPULATION IN 2012 – 2013. THE RESEARCH RESULTS OF THE ESSE-RF

Muromtseva G.A.¹, Kontsevaya A.V.¹, Konstantinov V.V.¹, Artamonova G.V.², Galaganova T.M.³,...

¹ FGBU State research center of preventive medicine of the Ministry of health of Russia, Moscow;

² FGBU Research Institute of complex problems of cardiovascular diseases SB RAMS, Kemerovo;

³ RD VPO North Ossetian state medical Academy, Vladikavkaz;..., Russia.

Information about the authors, where indicated:

full name, place of work of all authors, their positions, ORCID; full contact information is required for one (or more) of the author and includes e-mail, available phone number.

All members of the group of authors should meet all four criteria of authorship set forth in the ICMJE recommendations: 1) concept and design development or data analysis and interpretation, and 2) manuscript justification or verification of critical intellectual content, and 3) final approval for publication of the manuscript, and 4) consent to be responsible for all aspects of the work, and assume that issues relating to the thoroughness and diligent execution of any part of the study submitted are duly investigated and resolved. This information should also be contained in the document.

If the submitted material has authors who do not meet the criteria of authorship, but have made some contribution to the work, they should be listed in this

document and at the end of the article in the section of Acknowledgements.

Information on conflict of interest / funding.

The section contains the disclosure by all authors of possible relations with industrial and financial organizations that may lead to a conflict of interest in connection with the material presented in the manuscript. It is desirable to list the sources of funding for the work. If there is no conflict of interest, it is written: "Conflict of interest is not declared." Information on the existence of a conflict of interest should also be reflected in the Conflict of interest section at the end of the article.

Information about grants. Should be mentioned at the end of the article in the section Acknowledgements and at the end of the section Material and methods — with a full description of the role of the source of funding in the performance of work (design, information collection, analysis, data interpretation, etc.).

Information and ethics in the study.

Example of design:

The study was carried out in accordance with the standards of good clinical Practice (Good Clinical Practice) and the principles of the Helsinki Declaration. The study Protocol was approved by the Ethical committees of all participating clinical centers. Prior to being included in the study, written informed consent was obtained from all participants.

This information should also be reflected in the Material and methods section of the article.

All additional information (permits, questionnaires, etc.) can be requested from the authors in addition to the preparation of the work for printing.

Information on overlapping publications (if available).

Copyright. The use of any material (tables, figures) marked with a copyright icon in the article should be confirmed by a special permission from the author or publisher.

Information about the obtained consent in patients for the study.

Obtaining consent from patients for the study should also be reflected in the Material and methods.

For all clinical trials: information about the registration and placement of data on the study in any public register of clinical trials. The term "clinical study" refers to any research project that affects people (or groups of subjects) with/or without a comparative control group, studies the interaction between inter-

ventions to improve health or the results obtained. The world health organization offers the primary register: International Clinical Trials Registry Platform (ICTRP) (www.who.int/ictcp/network/primary/en/index.html). The clinical study is considered to be reliable in a group of more than 20 patients.

The number of words in the article (excluding summaries, sources of literature, figure captions and tables), the number of tables and figures.

The absence of an information file or incomplete text (not containing the above items) is the basis for refusal to accept the manuscript for consideration.

IV. Manuscript submission check-list

Since the main file of the manuscript is automatically sent to the reviewer for «blind review», it should not contain the names of the authors and institutions. The file contains only the following sections:

Article title

Summary with key words

List of abbreviations

Text

Acknowledgements (if any)

List of references

Tables, figures (if they can be embedded in the text of Word format).

The article title is written in capital letters (PREVALENCE of RISK FACTORS...), the end point is not needed. The title should clearly reflect the purpose of the work.

Summary with key words-sections are drawn up each with a separate line, highlighted in bold. The abstract should contain only those sections that are described in the rules for authors. For example, there is no section "Relevance" in the summary. The authors prescribe the relevance of their work in the introductory section of the manuscript.

List of abbreviations — when compiling a list of abbreviations to the article, including text, tables and figures, only those used by the author 3 or more times are included. Usually shrink often used in manuscripts of the terms (e.g., hypertension, CHF FC) and title of clinical trials (SOLVD, TIMI, HOPE).

The first reference to an abbreviation is always accompanied by the full spelling of the abbreviated concept, and the abbreviation is indicated in brackets. For example, blood pressure (BP); heart rate (HR). Capital letters are more often used to denote abbreviations. If abbreviations are used only in tables and

figures, and are not used in the text, they should not be included in the list of abbreviations, but should be given a transcript in the note to the table or figure. The summary of the article, as a separate document, is subject to the same rules as the article (abbreviations are made when they are used 3 or more times).

Abbreviations should be generally accepted and understandable to the reader, in accordance with the generally accepted norms in the scientific literature. Undesirable abbreviations that coincide in writing with others that have a different meaning.

Abbreviations in the list of abbreviations are written in alphabetical order, separated by commas, in solid text, using "dash". **Example of design:** BP-blood pressure, HR-heart rate.

Text — the text of the manuscript of the original works should be structured: Introduction, Material and methods, Results, Discussion and Conclusion. The text of reviews and lectures can be unstructured.

Text is printed on A4 sheet, font size — 12 pt, line spacing — 1.5, margins 2 cm on all sides. The system of SI units is used for processing the material, the % sign is put through a space from the number, the value of p is written with a semicolon: $p < 0.0001$; the value of n is written with a small letter ($n=20$); signs $>$, $<$, $\pm$, $=$, $+$, $-$ when numerical values are written without a space; the value of "year" or "year" is issued — 2014 or 2002 — 2014.

The article should be carefully verified by the author (s). The authors are responsible for the correctness of citation, doses and other factual materials.

Introduction — it is necessary to describe the context and prerequisites of the work (what is the essence of the problem and its significance). It sets certain goals or describes the object of the study, or a hypothesis that needs to be tested by comparison or observation. Only those sources that directly indicate the problem are cited.

Statistics — all published materials are reviewed by an expert in statistics and must meet "Uniform requirements for manuscripts submitted to biomedical journals" (Uniform Requirements for Manuscripts Submitted to Biomedical Journals, Ann Intern Med 1997, 126: 36 — 47). In the preparation of the statistical part of the work it is recommended to use special guidelines, for example, the European journal of cardiology: www.oxfordjournals.org/our_journals/eurheartj/for_authors/stat_guide.html

Statistical methods are described in detail in the Material and methods section.

Acknowledgements — all participants who do not meet the authorship criteria should be listed in the Acknowledgements section, which is located at the end of the article before the Literature section.

Making graphs, diagrams and drawings — tables and figures should provide the reader with visual information, be interesting and educational. They should be placed after the text of the article, as the reviewer and editor look at the manuscript as a whole. However, to print in the journal (at the stage of creating a layout) graphics, diagrams and drawings are required in electronic form in the formats "MS Excel", "Adobe Illustrator", "Corel Draw", "MS PowerPoint", photos with a resolution of at least 300 dpi.

The names of the graphs and figures, as well as notes to them should be placed under the figure/graph or placed at the end of the article.

These files are referred to as additional files. Figures should not repeat the materials of the tables.

Tables should contain the compressed, necessary data. Each table is placed at the end of the text (after the list of references) with the number, name and explanation (note, abbreviations).

The tables should clearly indicate the dimension of the indicators and the form of data ($M \pm m$; $M \pm SD$; Me ; Mo ; percentiles, etc.). All figures, totals and percentages should be carefully verified, and also correspond to the mention in the text. The explanatory notes are given below the table, if necessary. The footnotes must be in the following order: *, †, §, ||, ¶, #, **, †† etc.

Abbreviations should be listed in a footnote below the table in alphabetical order (for tables its list of abbreviations!).

Each first mention of a figure or table in the text is highlighted with a yellow marker. If a reference to a figure or table is included in the sentence, the full spelling of the word «figure 1», «table 1» is used; if the words are enclosed in brackets, the abbreviation is used (Fig. 1), (table. 1).

Providing the main file of the manuscript with the names of the authors or institutions is the basis for refusal to accept the manuscript for consideration.

V. The list of references.

In the form to fill in when submitting the article provides a list of cited literature (section — Literature).

Literary references are listed in the order of citation in the manuscript. The text refers to the serial number of the cited work in square brackets [1] or [1, 2]. Each link in the list is on a new line. All documents referred to in the text should be included in the list of references.

References to works that are not in the list of references and Vice versa, references to unpublished works, as well as to works of many years ago (>10 years) are not allowed. The only exceptions are rare highly informative works. Especially close attention to this item, please pay to those authors who submit "literature Review".

The bibliographic description contains the names of the authors up to three, after which, for domestic publications should indicate "et al.", for foreign — "et al." When citing articles from journals indicate in the following order the output: the name and initials of the authors, the name of the source, year, volume, number, pages (from and to). When citing articles from the collections indicate the output: name, initials, title, title of the collection, place of publication, year of publication, page (from and to).

If you want to make a quotation of the authors' names in the text, you must specify the name of the first author with the initials, the year of work. Example design: Smith AA, et al. (2018).

With the purpose of increase of citation in the journal is the transliteration of Russian sources with the use of the official languages in the following order: the authors and the journal title is transliterated in the Latin alphabet, and the name of the article is semantic transliteration (translation into English). The name of the source where the work is published is transliterated in Latin if the source (journal) does not have an official name in English).

All Russian-language sources of literature should be presented in the transliterated version of the model given below.

The author (s) are responsible for the correctness of the data given in the references.

The list of references should correspond to the format recommended by the American National organization For information standards (national Information Standards organization — NISO), adopted by the National Library of Medicine (NLM) for databases (Library's MEDLINE/PubMed database) NLM: <http://www.nlm.nih.gov/citingmedicine> Oh? The names of periodicals may be abbreviated. Usually

this form of writing is accepted by the publisher; it can be found on the website of the publisher, or in the list of abbreviations Index Medicus.

Mandatory all articles DOI specified, all books ISBN. References to dissertations, patents, theses and any collections without output and ISBN are not accepted.

Examples of link design:

Article citation:

Smith A, Jones B, Clements S. Clinical translation of tissue-engineered airway. *Lancet*. 2008;372:1201 — 09. DOI:10.00000/0000-0000-.

Russian-language sources with transliteration:

Bart BYa, Larina VN, Brodskyi MS, et al. Cardiac remodelling and clinical prognosis in patient with chronic heart failure and complete left bundle branch block. *Russ J Cardiol*. 2011;6:4 — 8. Russian. Барт Б. Я., Ларина В. Н., Бродский М. С., и др. Ремоделирование сердца и прогноз больных с хронической сердечной недостаточностью при наличии полной блокады левой ножки пучка Гиса. *Российский кардиологический журнал*. 2011;6:4 — 8. DOI:10.15829/1560-4071-2011-6-4-8.

Book:

Shlyakhto EV, Konradi AO, Tsyrlin VA. The autonomic nervous system and hypertension. SPb.: Meditsinskoe izdatel'stvo; 2008. Russian. Шляхто Е. В., Конради А. О., Цырлин В. А. Вегетативная нервная система и артериальная гипертензия. СПб.: Медицинское издательство; 2008. ISBN 0000 — 0000.

Chapter:

Nichols WW, O'Rourke MF. Aging, high blood pressure and disease in humans. In: Arnold E, ed. *McDonald's Blood Flow in Arteries: Theoretical, Experimental and Clinical Principles*. 3rd ed. London/Melbourne/Auckland: Lea and Febiger; 1990. p.398 — 420. ISBN 0000 — 0000.

Russian chapter:

Diagnostics and treatment of chronic heart failure. In: *National clinical guidelines 4th ed*. Moscow: Silicea-Polygraf; 2011. pp.203 — 93. Russian Диагностика и лечение хронической сердечной недостаточности. В кн: Национальные клинические рекомендации. 4-е издание. М.: Силицея-Полиграф; 2011.с.203 — 96. ISBN 0000 — 0000.

Webpage:

Panteghini M. Recommendations on use of biochemical markers in acute coronary syndrome:

IFCC proposals. eJIFCC 14. <http://www.ifcc.org/ejifcc/vol14no2/1402062003014n.htm> (28 May 2004)

All sources of literature are checked for correctness through the system of the Russian electronic library. Significant errors in citation or duplication of the source are the reason for the return of the manuscript to the authors for revision.

VI. Preparation of manuscript.

The author prepares the following documents to upload the manuscript to the site:

The main file is the text of the article (the system renames it after loading, so it does not matter how it is called).

Additional files-Directional (accompanying) letter, Information file with the Title page, information about the authors and disclosure of conflicts of interest, files with pictures.

For more information on placing articles on the website you can read <http://cardiovascular.elpub.ru/jour/announcement>

VII. Copyright and publishing policy.

This section regulates the relationship between the editorial Office (Publisher) of *International heart and vascular disease journal* (the "editorial Office") and the author or group of authors who submitted their manuscript for publication in the *International heart and vascular disease journal* (the "Author").

The author, by sending the article to the Editor, agrees that the editorial Board of the journal shall be transferred to the exclusive property rights to use the manuscript (transferred to the Editorial Board of the journal material, including such protected objects of copyright as photos of the author, drawings, diagrams, tables, etc.), including the reproduction in print and on the Internet; distribution; translation into any languages of the peoples of the world; export and import of copies of the journal with the article of the Author for distribution, to bring to the public.

The editorial Board reserves the right to reduce and edit the materials of the manuscript, to carry out scientific editing, to reduce and correct articles, to change the design of graphs, drawings and tables to bring into line with the design of the journal, without changing the meaning of the information provided.

When using the article, the editors have the right to supply it with any illustrated material, advertising and allow third parties to do so.

The editorial Board has the right to assign the rights received from the Author to third parties and has the right to prohibit third parties from any use of materials published in the journal for commercial purposes.

The author guarantees that he has exclusive rights to use the submitted material. In case of violation of this guarantee and the presentation of claims to the editorial Board, the Author independently and at his own expense undertakes to settle all claims. The editorial Board is not responsible to third parties for violation of the Author's guarantees.

The Author retains the right to use the published material, its fragments and parts for personal, including scientific and teaching purposes.

The Author transfers the above rights to the Editors without limitation of their validity period, in the territory of all countries of the world without limitation, including the territory of the Russian Federation.

The rights to the manuscript are considered to be transferred By the author of the editorial Office from the moment of sending an information letter about the acceptance of the manuscript to the press.

Reprinting of materials published in the journal by other individuals and legal entities is possible only with the written permission of the editorial Board, with the obligatory indication of the journal name, number and year of publication.

The editors are not responsible for the accuracy of the information provided by the Author.

The author, sending the manuscript to the Editor, gives permission to use and process personal data.

The editorial Board reserves the right to reduce and correct the articles, to change the design of graphs, figures and tables to comply with the standard of the journal, without changing the meaning of the information provided. In case of untimely response of the author (s) to the request of the editorial Board, the editorial Board may at its discretion make changes to the article or refuse to publish.

Sending to the editor of works that have already been sent to other publications or printed in them is absolutely not allowed. The editors are not responsible for the accuracy of the information provided by the authors. Articles sent in violation of the rules of registration are not accepted by the editorial Board for consideration.

VIII. The procedure for reviewing manuscripts

The manuscript should be sent in electronic form to the Editor through the website — [http://www.](http://www.heart-vdj.com)

[heart-vdj.com](http://www.heart-vdj.com). The manuscript should be drawn up in accordance with these requirements for scientific articles submitted for publication in the journal.

The author is sent a notification letter of receipt of the manuscript with the number (ID), which will be used in subsequent correspondence. The author can track the stages of work on his manuscript through the site. Since the process of bringing the manuscript to the necessary standards takes enough expert time, the payment for the initial review of the article was introduced, which the author (s) are required to carry out after the article is posted on the site.

The manuscript must pass the primary selection: the Editorial Board has the right to refuse publication or send comments to the article, which must be corrected by the Author before reviewing.

- checking the completeness of the manuscript: if you do not comply with the requirements of the Rules for the authors to complete the manuscript or its design, the Editors have the right to refuse to publish or in writing to require to send the missing materials or to correct the version already downloaded to the site.

- Manuscripts are checked in the "Antiplagiat" system. The originality of the manuscript should be at least 75 %. We expect manuscripts submitted for publication to be written in an original style that involves new thinking without the use of previously published text. Manuscript with originality below 75 % shall not be admissible.

All manuscripts submitted to the journal are sent to one of the permanent reviewers or an independent expert according to the profile of the research.

The review process is anonymous both for the Author and for the reviewers. The manuscript is sent to the reviewer without the names of the authors and the name of the institution.

The editorial Board informs the Author of the results of the review by e-mail.

If the reviewer makes a conclusion about the possibility of publication of the article and does not make significant corrections, the article is given to the expert on statistics and after a positive report is accepted for further work.

If the reviewer makes a conclusion about the possibility of publication of the article and gives instructions on the need for its correction, the Editorial Board sends the review to the Author with a proposal to take into account the recommendations of the reviewer in the preparation of a new version of the ar-

ticle or to refute them. In this case, the Author needs to make changes to the last version of the article file, which is located on the site (download file from the site, make changes and place the corrected article again, after removing the primary (uncorrected) version). The revised article is re-sent for review, and the conclusion is given that all the recommendations of the reviewer were taken into account. After receiving a positive response of the reviewer, the article is given to the expert on statistics and after a positive report is accepted for further work.

If the reviewer makes a conclusion about the impossibility of publication of the article. The author of the reviewed work is given the opportunity to read the text of the review, if he does not agree with the conclusions of the reviewer. In case of disagreement with the opinion of the reviewer, the Author has the right to provide a reasoned response to the Editor. The article can be sent for re-review or for approval to the editorial Board. The editorial Board or its authorized editor shall send its response to the Author.

All manuscripts that have been reviewed and evaluated by an expert in statistics are submitted to the editorial Board, which decides on the publication. After the decision on the admission of article for publication, the Editorial office inserts the publication of the article in terms of publications. Information about the annual (thematic) plan of publications is placed on the website of the journal.

The decision to publish a manuscript is made solely on the basis of its significance, originality, clarity of presentation and compliance of the research topic with the direction of the journal. Reports on studies in which negative results are obtained or the provisions of previously published articles are challenged are considered on General grounds.

Original reviews are kept in the Editorial office for 5 years from the date of publication.

In case of a decision to refuse to publish an article, its archive copy remains in the electronic system of the editorial Board, but access to it by editors or reviewers is closed.

IX. The manner of publication of manuscripts

According to the requirements of the Higher attestation Commission, the journal provides priority for post-graduate and doctoral works, the period of their publication depends on the expected date of protec-

tion, which the authors must specify in the primary documents attached to the manuscript.

Each issue of the journal is formed by a separate Executive editor appointed by the editor-in-Chief and/or editorial Board. It is the responsibility of the editor-in-charge to select high-quality articles for publication, and he can be guided by both thematic principles and a separate scientific direction.

All selected articles are submitted to the scientific editor and proofreader. After creating the layout of the article and editing it, the article will be available to the Author through the site. At this stage, it will be possible to send comments on the text of the article. The author is obliged to send his / her consent to the publication or his / her comments within the established time specified in the cover letter.

The editorial office does not send the author's copy by mail or PDF of the article by e-mail, access to the published numbers is open.

Subscription to the printed version is carried out by half a year (through subscription agencies).

X. After the publication in the journal

Information on publication is distributed in the following scientific citation databases: Russian science citation index, CYBERLENINKA and others. The article is assigned a DOI index and the full text is publicly available on the journal's website.

Information about the publication of the issue is distributed by mailing of The Cardioprogress Foundation and in social networks.

We expect the authors of the articles to actively make efforts to bring the results of their research to the public, namely: to have a personal page on the Internet (personal page), to monitor and update your profile ORCID and RecsearcherID, to involve colleagues in their work through social networks.

XI. Revocation or correction of articles

The full text of the journal's policy on Revocation and correction of articles is available in the information section on the website. The editors follow COPE Recommendations issued by the Committee on publishing ethics (COPE) — <http://www.publicationethics.org.uk>. in cases:

Editors of journals should consider the opinion of the publication, if:

they have clear evidence of the unreliability of the information published, either as a result of conscious actions (for example, falsification of data), or due to good faith errors (for example, errors in calculations or experiments); the findings have been previously published in another publication and there is no proper reference, authorization and justification for re-publication (i.e. duplicate publication.); it is plagiarism; describes unethical research.

Editors of journals should consider the concerns, if:

they received information about the authors' inappropriate actions, but there is no clear evidence of such behavior; there are arguments that the results of the work are unreliable, and the institution in which the authors work is not going to find out the truth; they believe that the investigation into the alleged violations committed by the authors in connection with the publication has either not been or will not be fair, impartial and convincing; the authors' violations are being investigated, but the results are not expected soon enough.

Journal editors should consider making amendments if:

a small part of the rest of the high-quality publication is unreliable (especially because of conscientious errors); the list of authors / sponsors contains errors (i.e., it does not contain someone who is worthy to be an author, or a person who does not meet the authorship criteria).

In most cases, a review is not appropriate if:

authorship needs to be changed, but there is no reason to doubt the validity of the findings.

XII. Position E-log backup (if journal is no longer published)

The purpose of backup is to prevent loss of information in case of hardware, software, critical and crisis situations, etc.

Information of the following main categories is subject to backup: — personal information of authors (personal directories on file servers); — pdf of published articles; — information about literary links to the article in the DOI system.

All this information is publicly available in The system of the Russian citation index on the website of the Electronic library www.elibrary.ru

XIII. Journal subscription

Information on subscriptions is available on the journal website in the section "Subscription":

XIV. Journal subscription

The name of the journal in English is International heart and vascular disease journal.

Official sites where information about the journal is placed:

<http://www.heart-vdj.com>

On the reception of the articles, making decisions about publication, reviews — mmamedov@mail.ru

On organizational issues (working with the site, subscription) — editor.ihvdj@gmail.com

Editorial office:

Room 213, Building 2, Prospect Gostinichny 6, Moscow 127106, Russia

e-mail: editor.ihvdj@gmail.com

Submission Preparation Checklist

As part of the submission process, authors are required to check off their submission's compliance with all of the following items, and submissions may be returned to authors that do not adhere to these guidelines.

The manuscripts are accepted if has not been published or submitted for publication elsewhere.

The file of the submitted article is in the format of a Microsoft Word document. It does not contain the names of the authors and institutions.

Files with a letter of transmittal and General information have been prepared for upload to the site.

3. The cited literature is presented in full, framed by the Rules for the authors and does not contain duplicates. All references are indicated in the text of the article.

Text should be typed with an interval of one line spacing, font Times New Roman, 12 pt; to highlight the accents it is recommended to use italics rather than underlining (except Internet links). All images, graphics and tables are placed within the text according to the meaning of the particular part of text (and not at the end of the document).

Text should follow the stylistic and bibliography requirements as stated in Regulations located in the Part "About Us."

Please, remove the authors' names from the title of the article and other parts of the document to ensure the anonymity of reviewing.

Copyright Notice

Authors who publish with this journal agree to the following terms:



Authors retain copyright and grant the journal right of first publication with the work simultaneously licensed under a Creative Commons Attribution License that allows others to share the work with an acknowledgement of the work's authorship and initial publication in this journal.

Authors are able to enter into separate, additional contractual arrangements for the non-exclusive distribution of the journal's published version of the work (e.g., post it to an institutional repository or publish it in a book), with an acknowledgement of its initial publication in this journal.

Authors are permitted and encouraged to post their work online (e.g., in institutional repositories or

on their website) prior to and during the submission process, as it can lead to productive exchanges, as well as earlier and greater citation of published work (See The Effect of Open Access).

Privacy Statement

Specified when registering the names and addresses will be used solely for technical purposes of a contact with the Author or reviewers (editors) when preparing the article for publication. Private data will not be shared with other individuals and organizations.

ISSN: 2309 — 0901 (Print)

ISSN: 2311 — 1631 (Online)

FOUNDATION FOR THE ADVANCEMENT OF CARDIOLOGY

“CARDIOPROGRESS”

knowledge, observation, action



The main functions of the Cardioprogress Foundation are:

- Research
- Education
- Science
- Publishing
- International collaboration
- Advertising and information

Official website: www.cardioproggress.ru

Tel: 007 965 236 1600

Email: inf.cardio@gmail.com

Moscow, Russia